

LINK BUNDLES AND INTERSECTION SPACES OF COMPLEX TORIC VARIETIES

MARKUS BANAGL AND SHAHRYAR GHAED SHARAF

ABSTRACT. There exist several homology theories for singular spaces that satisfy generalized Poincaré duality, including Goresky-MacPherson’s intersection homology, Cheeger’s L^2 cohomology and the homology of intersection spaces. The intersection homology and L^2 cohomology of toric varieties is known. Here, we compute the rational homology of intersection spaces of complex 3-dimensional toric varieties and compare it to intersection homology. To achieve this, we analyze cell structures and topological stratifications of these varieties and determine compatible structures on their singularity links. In particular, we compute the homology of links in 3-dimensional toric varieties. We find it convenient to use the concept of a rational homology stratification. It turns out that the intersection space homology of a toric variety, contrary to its intersection homology, is not combinatorially invariant and thus retains more refined information on the defining fan.

1. INTRODUCTION

A toric variety is a complex algebraic variety containing an algebraic torus as an open dense subset, such that the action of the torus on itself extends to the whole variety. Such varieties can be described by combinatorial data, called fans. Toric varieties are generally singular, and their homology does not usually satisfy Poincaré duality.

We focus here on compact complex 3-dimensional singular toric varieties associated to complete fans. For these, the fourth Betti number only depends on the number f_1 of 1-dimensional cones in the fan, but the second Betti number depends in addition on a parameter b , which then measures the discrepancy between these Betti numbers. The parameter b is not “combinatorial”, i.e. it depends on the precise form of the fan, not just on the number of cones in the various dimensions. If $b \neq 0$, then Poincaré duality is violated.

There are several cohomology theories available that restore duality. The intersection cohomology IH^* of Goresky and MacPherson [22], the L^2 -cohomology of Cheeger [12], [13], [14] for appropriately conical metrics on the regular part, and the homotopy-theoretic method of intersection spaces IX [6], yielding a cohomology theory $HI^*(X) := H^*(IX)$. Using a conical metric, Cheeger’s cohomology is isomorphic to intersection cohomology for a space with, say, only even-dimensional strata. While intersection homology of a complex 3-dimensional toric variety does repair Poincaré duality, it retains only a relatively small amount of the actual topology: The middle Betti number (indeed all odd Betti numbers) vanishes (Stanley [32]), and the second (and thus fourth) Betti number only depends on f_1 , as has been shown by Fieseler in [18].

It is well-known that IH^* and HI^* are generally not isomorphic. Does the homology of intersection spaces perhaps retain more information of the toric variety (while also repairing duality)? We find this indeed to be the case: One of our main results is the computation of HI^* for compact complex 3-dimensional toric varieties. We show that HI^* depends on f_1 , f_2 and b ; more precisely:

2020 *Mathematics Subject Classification.* 55N33, 14M25, 57S12, 57N80.

Key words and phrases. Toric Varieties, Intersection Homology, Stratified Spaces.

Theorem. (Theorem 68)

Let $X_{\mathcal{P}}$ be a compact complex 3-dimensional toric variety with m \mathbb{Q} -isolated singularities, $m \geq 1$, where \mathcal{P} is the underlying polytope. Let Σ be the dual fan to \mathcal{P} . We denote the number of 1-dimensional and 2-dimensional cones of Σ by f_1 and f_2 . Then

$$\begin{aligned} \mathrm{rk}(\tilde{H}_6(IX)) &= 0 \\ \mathrm{rk}(\tilde{H}_5(IX)) &= m - 1 \\ \mathrm{rk}(\tilde{H}_4(IX)) &= f_1 - 3 - b \\ \mathrm{rk}(\tilde{H}_3(IX)) &= 2(3f_1 - f_2 - b - 6) \\ \mathrm{rk}(\tilde{H}_2(IX)) &= f_1 - 3 - b \\ \mathrm{rk}(\tilde{H}_1(IX)) &= m - 1 \\ \mathrm{rk}(\tilde{H}_0(IX)) &= 0, \end{aligned}$$

where $b = \mathrm{rk}(H_4(X)) - \mathrm{rk}(H_2(X))$ and $\tilde{H}_*(-)$ denotes reduced singular homology.

Table 1 of the concluding Section 9 compares the above ranks to the ranks of intersection homology. The latter is obtained by imposing restrictions on how chains intersect the strata. Alternatively, one may wish to implement similar local modifications on the spatial level rather than the chain level. This motivates in part the theory of intersection spaces introduced by the first named author in [6]. To a stratified pseudomanifold X and perversity \bar{p} , one wishes to associate a space

$$I^{\bar{p}}X,$$

an *intersection space* of X , such that the ordinary reduced rational homology $\tilde{H}_*(I^{\bar{p}}X; \mathbb{Q})$ satisfies generalized Poincaré duality across complementary perversities when X is closed and oriented. In the absence of odd-co-dimensional strata and if \bar{p} is the middle perversity $\bar{p} = \bar{m}$, then $IX = I^{\bar{m}}X$ satisfies Poincaré self-duality rationally. The idea in forming intersection spaces is to use the homotopy cofiber of Moore approximations of singularity links. Thus we carry out a detailed analysis of these links in the toric case. An important structural feature that has been particularly emphasized by Agustín and de Bobadilla in their work on intersection space pairs [2] is that the link bundles of toric varieties can be trivialized, see Proposition 19 in the present paper. In [19], Fischli describes a procedure to endow compact toric varieties with CW structures. However, due to the resulting complexity in higher dimensions, the computation was carried out only up to dimension 2. We construct CW structures on complex 3-dimensional toric varieties and on their links. These structures allow us to determine the homology of all links, particularly of the real 5-dimensional ones (Proposition 33). This material may be of independent interest, notwithstanding the theory of intersection spaces. The CW structure on the links also allows us to construct their Moore approximations explicitly. We use a concept of \mathbb{Q} -homology stratified pseudomanifolds in order to be able to form the intersection space based on isolated singularity techniques. The observation here is that the real 3-dimensional links are rational homology spheres, so do not disturb duality. Homology stratifications are treated by Rourke and Sanderson in [28].

The theory HI^* has had applications in fiber bundle theory and computation of equivariant cohomology ([7]), K-theory ([6, Chapter 2.8],[31]), algebraic geometry (smooth deformation of singular varieties [10], perverse sheaves [8], mirror symmetry [6, Chapter 3.8]), theoretical Physics ([6, Chapter 3], [8]). An L^2 Hodge theorem for HI^* has been provided in [9] based on the use of the scattering metric. That theorem then provides a purely topological interpretation of spaces of extended weighted L^2 -harmonic forms with respect to fibered scattering metrics. A de Rham complex describing HI^* with \mathbb{R} coefficients was developed in [5] for stratification depth one and by Essig in [16] for higher stratification depth, see also [17]. Schlöder and Essig

obtained multiplicative de Rham theorems for intersection space cohomology in [29]. Higher stratification depth was addressed through the notion of intersection space pairs, introduced by Agustín Vicente and De Bobadilla in [2]. Verdier self-dual intersection space complexes are treated by Agustín Vicente, Essig and De Bobadilla in [1]. For isolated singularities, Klimczak [26] explained how to attach a top dimensional cell to IX so as to make it into a geometric Poincaré complex whose Poincaré duality isomorphism is given by capping with a fundamental class. Wrazidlo has generalized this to non-isolated singularities of depth one in [33].

2. PRELIMINARIES

Let us recall the definition of a rational polyhedral cone in \mathbb{R}^n .

Definition 1. We call $\sigma \subset \mathbb{R}^n$ a *rational polyhedral cone*, if there exist finitely many vectors $\nu_1, \dots, \nu_k \in \mathbb{Z}^n$ such that

$$(1) \quad \sigma = \{x_1\nu_1 + \dots + x_k\nu_k \mid x_1, \dots, x_k \in \mathbb{R}_{\geq 0}\}.$$

σ is **proper** if it is not spanned by any proper subset of $\{\nu_1, \dots, \nu_k\}$, and ν_1, \dots, ν_k lie strictly on one side of some hyperplane in \mathbb{R}^n .

The dimension of σ is defined to be the dimension of $\text{span}(\sigma) \subset \mathbb{R}^n$. Note also that σ is called simplicial if the generating vectors $\nu_1, \dots, \nu_k \in \mathbb{Z}^n$ can be chosen linearly independent. This implies that $k = \dim(\sigma)$.

Definition 2. We call a cone τ a (proper) *face* of cone σ if it is spanned by a (proper) subset of $\{\nu_1, \dots, \nu_k\}$ and contained in the topological boundary of σ . In this case, we write $\tau \preceq \sigma$ ($\tau \prec \sigma$ if τ is a proper face of σ).

Definition 3 (Complete rational polyhedral cone). A **complete rational polyhedral cone complex** (**complete fan**) is a nonempty set Σ of proper rational polyhedral cones in \mathbb{R}^n satisfying the following conditions:

- (1) If τ is a face of a cone $\sigma \in \Sigma$, then $\tau \in \Sigma$.
- (2) If $\sigma, \sigma' \in \Sigma$, then $\sigma \cap \sigma'$ is a face of both σ and σ' .
- (3) $\bigcup_{\sigma \in \Sigma} \sigma = \mathbb{R}^n$.

Note that $\{0\}$ is in Σ as Σ contains at least one cone and $\{0\}$ is a face on any cone.

Definition 4. Let Σ be a complete rational polyhedral cone complex in \mathbb{R}^n . Reversing the inclusions and the dimensional grading in the face lattice of Σ , we obtain the face lattice of an abstract polyhedron which can be realized in \mathbb{R}^n as a regular (polyhedral) cell complex. We denote this abstract polyhedron by $\mathcal{P}(\Sigma)$ and its geometrical realization by $|\mathcal{P}(\Sigma)|$. The abstract polyhedron $\mathcal{P}(\Sigma)$ is called **the dual polyhedron** or **the dual polytope**. If the fan Σ is understood we will simply write \mathcal{P} . By abuse of notation we will frequently write \mathcal{P} for $|\mathcal{P}|$. The dimension of $\mathcal{P}(\Sigma)$ is n , since the top dimensional cell of $\mathcal{P}(\Sigma)$ is associated to the cone $\{0\}$ and it is n -dimensional.

Remark 5. The faces in $\mathcal{P}(\Sigma)$ are in one-to-one correspondence (in complementary dimension) with the cones in Σ . This defines a bijection $\delta : \mathcal{P} \rightarrow \Sigma$ and we denote the dual cone to $\tau \in \mathcal{P}$ by $\delta(\tau) \in \Sigma$. In particular Σ can be reconstructed from \mathcal{P} . From that perspective we shall also write $\Sigma = \Sigma_{\mathcal{P}}$.

Definition 6. A CW complex is called **regular** if its characteristic maps can be chosen to be embeddings.

Definition 7. The 0-dimensional faces of \mathcal{P} are called **vertices**. The 1-dimensional faces are called **edges** and **facets** are faces with co-dimension 1. The set

$$\mathcal{P}^i = \{F : F \text{ is a face of } \mathcal{P}, \dim(F) \leq i\}$$

is the i -skeleton of \mathcal{P} .

Regular CW structure of the dual polytope \mathcal{P} .

Note that \mathcal{P} is homeomorphic to an n -disc \mathcal{D}^n , where $\dim(\mathcal{P}) = n$. The structure of the underlying fan induces a regular CW structure on \mathcal{P} as follows:

The vertex $\{0\} \in \Sigma$ is dual to the interior of \mathcal{P} , represented by an n -dimensional cell in the CW structure. Each k -dimensional cone in Σ is dual to an $(n-k)$ -dimensional face of \mathcal{P} , represented by an $(n-k)$ -dimensional cell in the CW structure. Note also that the boundary maps are induced from the inclusion of faces in Σ (or dually in \mathcal{P}). This gives us a CW complex on $\mathcal{P} \cong \mathcal{D}^n$, which we will use later. Although we do not require the exact form of the boundary operators of \mathcal{P} in our preceding discussion, we briefly describe the attaching maps. As usual, we start with a discrete set, X^0 , whose points represent the vertices of \mathcal{P} . We glue each 1-dimensional cell to its topological boundary in \mathcal{P} , which consists of its neighboring 0-dimensional cells. Inductively, we attach each k -cell, representing a k -dimensional face of \mathcal{P} , to its lower-dimensional neighboring faces. Note that $|\mathcal{P}| - \text{int}(\mathcal{P}) \cong \mathcal{S}^{n-1}$. In the last step we attach $\text{int}(\mathcal{P})$, represented by the n -cell, to its topological boundary $|\mathcal{P}| - \text{int}(\mathcal{P})$. Note that due to the fact that each k -dimensional face of \mathcal{P} is homeomorphic to \mathcal{D}^k , each characteristic map can be chosen to be an embedding. Thus, the above CW structure is a regular CW structure.

Let $T^n = \mathbb{R}^n / \mathbb{Z}^n \cong \overbrace{\mathcal{S}^1 \times \cdots \times \mathcal{S}^1}^n$ be an n -torus. The projection map $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^n / \mathbb{Z}^n$ maps a **rational** k -dimensional linear subspace $\mathbf{V} \subset \mathbb{R}^n$ to a compact subtorus $\pi(\mathbf{V}) \subset T^n$. The **rationality** here means that \mathbf{V} has a basis in \mathbb{Z}^n . Now each affine k -plane parallel to \mathbf{V} in \mathbb{R}^n determines a subtorus “parallel” to $\pi(\mathbf{V})$. Collapsing each of these parallel subtori to a point will give us a compact subspace $T^n / \pi(\mathbf{V}) \subset T^n$, which is obviously homeomorphic to T^{n-k} .

Toric varieties.

In this work, we mainly study compact toric varieties, from a topological perspective. Hence, we restrict ourselves to toric varieties associated with complete fans. This ensures that the resulting toric variety is compact. Compact toric varieties are even-dimensional topological pseudomanifolds with only even dimensional strata. We recall the description of toric varieties as given by Yavin in [35].

Topologically, we can construct the toric variety associated with a **complete rational cone complex** Σ (**complete fan**) as follows:

Let \mathcal{P} be the dual polytope of Σ and $\bar{X} = \mathcal{P} \times T^n$. Each $\sigma_D = \delta(\sigma)$, a k -dimensional face of \mathcal{P} , is dual to an $(n-k)$ -dimensional cone σ in Σ . Let \mathbf{V}_σ be the linear span of σ in \mathbb{R}^n . Now if $x \in \text{int}(\sigma_D)$ we collapse $\{x\} \times T^n$ to $\{x\} \times T^n / \pi(\mathbf{V}_\sigma)$. This procedure can be done for each face of \mathcal{P} and the resulting space $X_{\mathcal{P}}$ or (X_Σ) is called the **toric variety** associated with Σ (or \mathcal{P}). Note that for each $x \in \mathcal{P}$ there is a unique σ_D such that $x \in \text{int}(\sigma_D)$. Thus the above construction is well-defined.

Example 8 ($n = 2$). Let $n, m \in \mathbb{N}_{\geq 0}$, $(n, m) \neq 0$ and n and m be relatively prime. Consider the following fan:

$$\Sigma_2 = \{\{0\}, \tau_1, \tau_2, \tau_3, \tau_4, \sigma_{12}, \sigma_{23}, \sigma_{34}, \sigma_{41}\}$$

where

$$\tau_1 = \{x(n, m) \mid x \in \mathbb{R}_{\geq 0}\},$$


$$\tau_2 = \{x(-m, n) \mid x \in \mathbb{R}_{\geq 0}\},$$

$$\tau_3 = \{x(-n, -m) \mid x \in \mathbb{R}_{\geq 0}\},$$


$$\tau_4 = \{x(m, -n) \mid x \in \mathbb{R}_{\geq 0}\},$$

$$\sigma_{12} = \{x(n, m) + y(-m, n) \mid x, y \in \mathbb{R}_{\geq 0}\};$$



see Figure 1.

FIGURE 1. The 2-dimensional complete fan Σ_2 .

The cones σ_{23} , σ_{34} and σ_{41} are defined similarly. The dual polytope is homeomorphic to an 2-dimensional convex polygon with four 1-dimensional faces or simply a square. Thus we can write $\bar{X} = \mathcal{I} \times \mathcal{I} \times \mathcal{T}^2$. The torus \mathcal{T}^2 can be written as $\pi(\mathbf{V}_{\tau_1}) \times \pi(\mathbf{V}_{\tau_2})$. Hence \bar{X} can be rewritten as $(\mathcal{I} \times \pi(\mathbf{V}_{\tau_1})) \times (\mathcal{I} \times \pi(\mathbf{V}_{\tau_2}))$. Note that due to the structure of Σ_2 applying the topological construction of toric varieties yields that either $\pi(\mathbf{V}_{\tau_1})$ or $\pi(\mathbf{V}_{\tau_2})$ is subject to collapse on 1-dimensional faces of $\mathcal{P}(\Sigma_2)$. Based on this observation it is easy to show that $X_{\Sigma_2} \cong X_{\Sigma_1} \times X_{\Sigma_1} \cong \mathbb{P}^1 \times \mathbb{P}^1$, where $\Sigma_1 = \{\{0\}, \tau_1, \tau_3\}$; see Figure 2.

FIGURE 2. The toric variety associated to Σ_2 is homeomorphic to $\mathcal{S}^2 \times \mathcal{S}^2$.

Example 9 ($n = 3$). Consider the following fan, Σ_3 , where τ_z is generated by $\hat{i}_z = (0, 0, 1)$. The cones τ_x and τ_y are generated by \hat{i}_x and \hat{i}_y , respectively. The fourth 1-dimensional cone of Σ_3 is generated by $i_4 = (-1, -1, -1)$. The generators of the 2-dimensional cone σ_{xz} are \hat{i}_x and \hat{i}_z and the five remaining 2-dimensional cones are constructed in a similar manner. Finally, ω_{xyz} is generated by \hat{i}_x , \hat{i}_y and \hat{i}_z . The three remaining 3-dimensional cones are constructed similarly; see Figure 3. It is easy to show that the dual polytope is a tetrahedron. Applying the topological construction of toric varieties on Σ_3 yields a toric variety which is homeomorphic to \mathbb{P}^3 (Figure 4).

FIGURE 3. The complete fan Σ_3 in \mathbb{R}^3 .FIGURE 4. The toric variety X_{Σ_3} is homeomorphic to \mathbb{P}^3 .

3. TOPOLOGY OF TORIC VARIETIES

In this section, we study toric varieties from the perspective of stratified pseudomanifolds. First of all, we will recall that toric varieties are indeed pseudomanifolds. In the next step, we will verify that the link bundles of toric varieties are topologically trivial. Toric varieties are normal pseudomanifolds. In other words, the link of the stratum with depth 1 is always homeomorphic to a circle \mathcal{S}^1 .

Definition 10. We define a **topologically stratified space** inductively on dimension. A 0-dimensional topologically stratified space X is a countable set with the discrete topology. For

$m > 0$ an m -dimensional topologically stratified space is a para-compact Hausdorff topological space X equipped with a filtration

$$X = X_m \supseteq X_{m-1} \supseteq \cdots \supseteq X_1 \supseteq X_0 \supseteq X_{-1} = \emptyset$$

by closed subsets X_j such that if $x \in X_j - X_{j-1}$ there exists a neighborhood \mathcal{N}_x of x in X , a compact $(m - j - 1)$ -dimensional topologically stratified space \mathcal{L} with filtration

$$\mathcal{L} = \mathcal{L}_{m-j-1} \supseteq \cdots \supseteq \mathcal{L}_1 \supseteq \mathcal{L}_0 \supseteq \mathcal{L}_{-1} = \emptyset,$$

and a homeomorphism $\phi : \mathcal{N}_x \rightarrow \mathbb{R}^j \times \mathcal{C}(\mathcal{L})$, where $\mathcal{C}(\mathcal{L})$ is the open cone on \mathcal{L} , such that ϕ takes $\mathcal{N}_x \cap X_{j+i+1}$ homeomorphically onto

$$\mathbb{R}^j \times \mathcal{C}(\mathcal{L}_i) \subseteq \mathbb{R}^j \times \mathcal{C}(\mathcal{L})$$

for $m - j - 1 \geq i \geq 0$, and ϕ takes $\mathcal{N}_x \cap X_j$ homeomorphically onto

$$\mathbb{R}^j \times \{\text{vertex of } \mathcal{C}(\mathcal{L})\}.$$

Remark 11. It follows that $X_j - X_{j-1}$ is a j -dimensional topological manifold. (The empty set is a manifold of any dimension.) We call the connected components of these manifolds the **strata** of X . Any \mathcal{L} that satisfies the above properties is referred to as a **link** of the stratum at x .

Definition 12. An m -dimensional topological pseudomanifold is a para-compact Hausdorff topological space X which possesses a topological stratification such that $X_{m-1} = X_{m-2}$ and $X - X_{m-1}$ is dense in X .

Definition 13. We call a stratum (homologically) **singular** if none of its links is a homology sphere. A stratum is (homologically) **rationally singular** if none of its links is a rational homology sphere.

Now, we recall that there is a natural stratification of toric varieties. We also give an algebraic description of the situation.

Let X be a toric variety over a polytope \mathcal{P} . Consider the map $X \xrightarrow{p} \mathcal{P}$ introduced earlier. The preimage $X_{2i} = p^{-1}(\mathcal{P}^i)$ is a $2i$ -dimensional topological space. We claim that the filtration

$$(2) \quad X_{\mathcal{P}} = X_{2m} \supset X_{2(m-1)} \supset \cdots \supset X_2 \supset X_0,$$

is a stratification of the toric variety $X_{\mathcal{P}}$. The local homeomorphisms will be defined later. But first we need to give a topological description of links.

Construction of links

Let τ be a face of \mathcal{P} . Let \mathcal{M}_{τ} be an abstract polytope, geometrically realized as a subspace of \mathcal{P} , and defined as follows:

Let $\mathcal{S}_{\tau} = \{\sigma \in \mathcal{P}, \sigma \cap \tau \neq \emptyset \text{ and } \dim(\sigma) > \dim(\tau)\}$ be the set of all higher dimensional neighboring faces of τ in \mathcal{P} .

We shall define an abstract polytope \mathcal{M}_{τ} with the following properties:

- (1) $\forall \sigma \in \mathcal{S}_{\tau} : \exists ! \gamma_{\sigma} \in \mathcal{M}_{\tau}$ such that $\text{int}(\gamma_{\sigma}) \cap \text{int}(\sigma) \neq \emptyset$ with

$$\dim(\gamma_{\sigma}) = \dim(\sigma) - (1 + \dim(\tau))$$

and $\text{int}(\gamma_{\sigma}) \cap \text{int}(\sigma') = \emptyset$ if $\sigma' \in \mathcal{P}$ and $\sigma' \neq \sigma$.

- (2) If $\omega, \sigma \in \mathcal{S}_{\tau}$ with $\omega \prec \sigma$, then we demand that $\gamma_{\omega} \prec \gamma_{\sigma}$ and $\gamma_{\sigma} \cap \omega = \gamma_{\omega}$.

Note that $\text{int}(\sigma) \cong \text{int}(\mathcal{D}^{\dim(\sigma)})$. So, the first requirement can be satisfied by embedding $\text{int}(\mathcal{D}^{\dim(\gamma_{\sigma})})$ in $\text{int}(\mathcal{D}^{\dim(\sigma)})$. Yet the second condition describes how the boundary of $\mathcal{D}^{\dim(\gamma_{\sigma})}$ meets the boundary of $\mathcal{D}^{\dim(\sigma)}$. We refer to \mathcal{M}_{τ} as a **base** of the link of $x \in p^{-1}(\text{int}(\tau)) \subset X_{\mathcal{P}}$.

Proposition 14. *Let Σ be a complete fan in \mathbb{R}^n and \mathcal{P} its dual polytope. Given any face $\tau \in \mathcal{P}$ there exists an abstract polytope \mathcal{M}_τ satisfying the above conditions (1) and (2) and having a geometrical realization $|\mathcal{M}_\tau|$ as a subpolyhedron of a geometrical realization of \mathcal{P} .*

Proof. First, we show the existence of \mathcal{M}_τ . Let $\Sigma_{\mathcal{P}}$ be a complete fan and \mathcal{P} be the dual polytope associated with $\Sigma_{\mathcal{P}}$. Let $\delta : \mathcal{P} \rightarrow \Sigma_{\mathcal{P}}$ be the bijection defined in Remark 5. Let $\Sigma_{\mathcal{S}_\tau}$ be the set of all lower dimensional neighboring cones of $\tau \in \mathcal{P}$ in $\Sigma_{\mathcal{P}}$. Dual to the previous construction, we define the dual fan associated with \mathcal{M}_τ to be a fan in $\mathbb{R}^{\dim(\delta(\tau))-1}$ with the following property:

- $\forall \delta(\sigma) \in \Sigma_{\mathcal{S}_\tau} : \exists ! \delta(\gamma_\sigma) \in \Sigma_{\mathcal{M}_\tau}$ such that $\delta(\gamma_\sigma) \subset \delta(\tau)$ as a cone. In other words, $\delta(\gamma_\sigma)$ lies in the topological boundary of the cone $\delta(\tau)$.

One can verify easily that both sets of conditions are dually equivalent. In other words, there is an order-reversing bijection between $\Sigma_{\mathcal{S}_\tau}$ and \mathcal{S}_τ . The bijection is the restriction of δ^{-1} to $\Sigma_{\mathcal{S}_\tau}$. The set $\Sigma_{\mathcal{M}_\tau}$ is a set of cones such that

$$0 \leq \dim(\delta(\gamma_\sigma)) < \dim(\delta(\tau))$$

Hence, $\Sigma_{\mathcal{S}_\tau}$ is an $(\dim(\delta(\tau)) - 1)$ -dimensional cone-complex, embedded in $\mathbb{R}^{\dim(\delta(\tau))}$, a rational subspace of $\mathbb{R}^{\dim(\Sigma_{\mathcal{P}})}$. By cone-complex, we mean that $\Sigma_{\mathcal{M}_\tau}$ satisfies the conditions (1) and (2) of Definition 3. However, instead of the third axiom, we have

$$\bigcup_{\delta(\gamma_\sigma) \in \Sigma_{\mathcal{M}_\tau}} \delta(\gamma_\sigma) \cong \mathbb{R}^{\dim(\delta(\tau))-1}.$$

But the crucial point to bear in mind is the following. Reversing the inclusions and grading in $\Sigma_{\mathcal{M}_\tau}$ will still result in an abstract polytope, which we can geometrically realize in $\mathbb{R}^{\dim(\delta(\tau)-1)}$. Existence of the topological boundary of a cone yields the existence of \mathcal{M}_τ .

Now, we embed $\mathcal{C}(\mathcal{M}_\tau)$ in \mathcal{P} , geometrically. For a given point $x \in \text{int}(\tau)$, where $\dim(\tau) \geq 1$, let $\mathcal{V} \cong \mathbb{R}^{\dim(|\mathcal{M}_\tau|)+1}$ be an affine subspace of $\mathbb{R}^{\dim(|\mathcal{P}|)}$ which is orthogonal to τ in $\mathbb{R}^{\dim(|\mathcal{P}|)}$ and $x \in \mathcal{V}$. Note that $\dim(|\mathcal{M}_\tau|) = \dim(|\mathcal{P}|) - (1 + \dim(\tau))$, hence, such \mathcal{V} can always be found. Thus, we have $\sigma_\gamma \cap \mathcal{V} \neq \emptyset$ for each $\gamma \in \mathcal{M}_\tau$, where $\sigma_\gamma \in \mathcal{S}_\tau$ with $\text{int}(\gamma) \cap \text{int}(\sigma_\gamma) \neq \emptyset$. Now, we choose a geometrical realization of γ such that $\gamma \subset \sigma_\gamma \cap \mathcal{V}$. Note that $\dim(\sigma_\gamma \cap \mathcal{V}) = \dim(\gamma) + 1$. Thus, we can find such a geometrical realization consistently. Now, let $\mathcal{C}(|\mathcal{M}_\tau|) = (|\mathcal{M}_\tau| \times \mathcal{I}) / |\mathcal{M}_\tau| \times \{1\}$ be the cone of $|\mathcal{M}_\tau|$. We embed $\mathcal{C}(|\mathcal{M}_\tau|)$ into $|\mathcal{P}|$ as follows:

Let $v \in \mathcal{C}(|\mathcal{M}_\tau|)$ be the vertex of $\mathcal{C}(|\mathcal{M}_\tau|)$ and $\theta : \mathcal{C}(|\mathcal{M}_\tau|) \rightarrow |\mathcal{P}|$ a map such that for each $\gamma \in \mathcal{M}_\tau$,

$$\begin{aligned} \theta(\text{int}(\gamma) \times [0, 1]) &\subset \text{int}(\sigma_\gamma) \cap \mathcal{V}, \\ \theta(\gamma \times [0, 1]) &\subset \left(\text{int}(\sigma_\gamma) \cup \left(\bigcup_{\substack{\omega_\gamma \in \mathcal{S}_\tau \\ \omega_\gamma \prec \sigma_\gamma}} \text{int}(\omega_\gamma) \right) \right) \cap \mathcal{V} \\ \theta(\gamma \times \{0\}) &\cong |\mathcal{M}_\tau| \cap \text{int}(\sigma_\gamma) \cong \text{int}(\gamma). \end{aligned}$$

Note that for each $\eta, \gamma \in \mathcal{M}_\tau$ if $\gamma \prec \eta$ then we have $\sigma_\gamma \prec \sigma_\eta$ in \mathcal{S}_τ . We require that

$$(3) \quad \theta(\eta \times [0, 1]) \cap \sigma_\gamma = \theta(\gamma \times [0, 1]).$$

As the last requirement, we want $\theta(\text{int}(\gamma) \times [0, 1]) \hookrightarrow \text{int}(\sigma_\gamma)$ to be a topological embedding. Note that this can always be fulfilled because $\gamma \cong \mathcal{D}^{\dim(\gamma)}$ and $\sigma_\gamma \cong \mathcal{D}^{\dim(\sigma_\gamma)}$. At last, one should bear in mind that Equation (3) ensures that θ is also an embedding on the topological boundary of $\sigma_\gamma \cap \mathcal{V}$ for each $\sigma_\gamma \in \mathcal{S}_\tau$. We set $\theta(v) = x$. Note that because of $\gamma \subset \sigma_\gamma \cap \mathcal{V}$ and the fact that we can choose $\theta(\gamma \times [0, 1]) \hookrightarrow \text{int}(\sigma_\gamma)$ as an embedding, it is possible to choose θ continuous on $\mathcal{C}(|\mathcal{M}_\tau|)$ and thus an embedding of $\mathcal{C}(|\mathcal{M}_\tau|)$ into $|\mathcal{P}|$. Now let $\mathcal{V}^\perp \subset \mathbb{R}^{\dim(|\mathcal{P}|)}$ be

an orthogonal affine subspace to \mathcal{V} such that $\tau \subset \mathcal{V}^\perp$. Choose \mathcal{U} a neighborhood of x in $|\mathcal{P}|$ such that $\mathcal{U} \cap \mathcal{V}^\perp \cong \text{int}(\mathcal{D}^{\dim(\tau)}) \cong \mathbb{R}^{\dim(\tau)} \cong \text{int}(\tau)$ and $\mathcal{U} \cap \mathcal{V} \cong \mathcal{C}(|\mathcal{M}_\tau|)$. Thus, we have

$$(4) \quad \mathcal{U} \cong \mathbb{R}^{\dim(\tau)} \times \mathcal{C}(|\mathcal{M}_\tau|).$$

□

Remark 15. Consider \mathcal{M}_τ and \mathcal{M}'_τ such that they both satisfy the above conditions. Then for each $\sigma \in \mathcal{S}_\tau$ and $\gamma_\sigma \in \mathcal{M}_\tau$ there exists a $\gamma'_\sigma \in \mathcal{M}'_\tau$ with $\dim(\gamma_\sigma) = \dim(\gamma'_\sigma)$. Note that the construction of \mathcal{S}_τ and the uniqueness in the first condition implies that $\text{card}(\mathcal{M}'_\tau - \mathcal{M}_\tau^{i-1})$ i.e. the number of i -dimensional faces of \mathcal{M}'_τ is equal to $\text{card}(\mathcal{M}'_\tau^i - \mathcal{M}_\tau^{i-1})$. Thus, there is a bijection between \mathcal{M}_τ and \mathcal{M}'_τ .

Now, let $|\mathcal{M}_\tau| \subset |\mathcal{P}|$ be a geometrical realization of \mathcal{M}_τ and $n = \dim(|\mathcal{P}|)$. Recall the bijection $\delta : \mathcal{P} \rightarrow \Sigma_{\mathcal{P}}$ that we introduced earlier. For each $\gamma \in \mathcal{M}_\tau$ choose $\sigma_\gamma \in \mathcal{S}_\tau$ such that $\gamma \cap \sigma_\gamma \neq \emptyset$. Note $\tau \prec \sigma_\gamma$ and thus $\delta(\sigma_\gamma) \prec \delta(\tau)$ in $\Sigma_{\mathcal{P}}$. This implies that $\pi(\delta(\sigma_\gamma)) \subset \pi(\delta(\tau))$ in T^n . We construct \mathcal{L}_τ , the link of a point $x \in p^{-1}(\text{int}(\tau))$, by means of the following map:

$$(5) \quad \begin{aligned} \mathcal{M}_\tau \times T^n &\longrightarrow \mathcal{L}_\tau \\ \{y\} \times T^n &\longmapsto \{y\} \times T^n / \left(\pi(\delta(\sigma_\gamma)) \times (T^n / \pi(\delta(\tau))) \right), \end{aligned}$$

where $y \in \text{int}(\gamma)$. Note that similar to the topological construction of toric varieties,

$$T^n / \left(\pi(\delta(\sigma_\gamma)) \times (T^n / \pi(\delta(\tau))) \right)$$

is defined by collapsing $\pi(\delta(\sigma_\gamma))$ and each parallel torus to $\pi(\delta(\sigma_\gamma))$ in T^n to a point and then collapsing each parallel torus to $T^n / \pi(\delta(\tau))$ in $T^n / \pi(\delta(\sigma_\gamma))$ to a point. Due to the previous consideration, $T^n / \left(\pi(\delta(\sigma_\gamma)) \times (T^n / \pi(\delta(\tau))) \right)$ is well-defined.

Remark 16. Note that for each $y \in \text{int}(\gamma)$, $T^n / \left(\pi(\delta(\sigma_\gamma)) \times (T^n / \pi(\delta(\tau))) \right) \subseteq T^n / \pi(\delta(\sigma_\gamma))$ and hence $\mathcal{L}_\tau \subset X$, where $n = \dim(|\mathcal{P}|)$.

Remark 17. The dimension relation $\dim(|\mathcal{M}_\tau|) = \dim(|\mathcal{P}|) - (1 + \dim(\tau))$ holds. This comes from the fact that $\text{int}(|\mathcal{P}|)$ considered as a face in \mathcal{P} , which is dual to $\{0\} \in \Sigma_{\mathcal{P}}$, is a higher dimensional neighboring face of each $\tau \in \mathcal{P}$. So $\dim(\mathcal{L}_\tau) = \dim(|\mathcal{P}|) - (1 + \dim(\tau)) + \dim(\pi(\delta(\tau)))$. Note that $\dim(\tau) = \dim(|\mathcal{P}|) - \dim(\delta(\tau))$. Thus, we have

$$\dim(\mathcal{L}_\tau) = \dim(X_{\mathcal{P}}) - (1 + 2 \dim(\tau)).$$

Remark 18. There is a natural projection $\mathcal{L}_\tau \xrightarrow{p_{\mathcal{L}}} \mathcal{M}_\tau$. Similarly, we define

$$(\mathcal{L}_\tau)_{2i+1} = p_{\mathcal{L}}^{-1}(\mathcal{M}_\tau^i).$$

At this point, we want to recall that there is a natural stratification of toric varieties. We will also give an algebraic description of the strata. We claim that the filtration given in Expression (2), is a stratification in the sense of Definition 10.

Proposition 19. Let Σ be a complete fan and \mathcal{P} the associated dual polytope. Then $X_{\mathcal{P}}$, the associated toric varieties with \mathcal{P} , is a topological pseudomanifold with a trivial link bundle.

Proof. We can construct \mathcal{U} in the proof of Proposition 14 as follows. We embed $\mathcal{C}(|\mathcal{M}_\tau|)$ in \mathcal{V} as in that proof. Now, we embed $\mathring{\mathcal{D}}^{\dim(\tau)} \times \mathcal{C}(|\mathcal{M}_\tau|)$ in $\mathcal{V} \times \mathcal{V}^\perp \cong \mathbb{R}^{\dim(|\mathcal{P}|)}$ such that

$$(\mathring{\mathcal{D}}^{\dim(\tau)} \times \mathcal{C}(|\mathcal{M}_\tau|)) \cap \mathcal{V}^\perp = \mathring{\mathcal{D}}^{\dim(\tau)}.$$

Finally, we set $\mathcal{U} \cong \mathring{\mathcal{D}}^{\dim(\tau)} \times \mathcal{C}(|\mathcal{M}_\tau|) \cong \mathbb{R}^{\dim(\tau)} \times \mathcal{C}(|\mathcal{M}_\tau|)$.

For each $\sigma_\gamma \in \mathcal{S}_\tau$, we have $\mathcal{U} \cap \text{int}(\sigma_\gamma) \cong (\mathbb{R}^{\dim(\tau)} \cap \text{int}(\sigma_\gamma)) \times (\mathcal{C}(|\mathcal{M}_\tau|) \cap \text{int}(\sigma_\gamma))$. Keep also in mind that $(\bigcup_{\gamma \in \mathcal{M}_\tau} \text{int}(\sigma_\gamma)) \cup \text{int}(\tau)$ is an open cover of \mathcal{U} . Thus

$$p^{-1}(\mathcal{U}) = \bigcup_{\gamma \in \mathcal{M}_\tau} \left((\text{int}(\sigma_\gamma) \cap \mathcal{U}) \times T^n / \pi(\delta(\sigma_\gamma)) \right) \cup \left((\text{int}(\tau) \cap \mathcal{U}) \times T^n / \pi(\delta(\tau)) \right).$$

Recall that $\tau \prec \sigma_\gamma$ in \mathcal{P} , hence $\delta(\sigma_\gamma) \prec \delta(\tau)$ which implies $\pi(\delta(\sigma_\gamma)) \subset \pi(\delta(\tau))$ and $T^n / \pi(\delta(\tau)) \subset T^n / \pi(\delta(\sigma_\gamma))$. Now, define $T_\tau = T^n / \pi(\delta(\tau))$. Then, we have

$$p^{-1}(\mathcal{U}) = T_\tau \times \left(\bigcup_{\gamma \in \mathcal{M}_\tau} \left((\text{int}(\sigma_\gamma) \cap \mathcal{U}) \times T^n / (\pi(\delta(\sigma_\gamma)) \times T_\tau) \right) \cup (\text{int}(\tau) \cap \mathcal{U}) \right).$$

Using Homeomorphism (4), we arrive at

$$p^{-1}(\mathcal{U}) = (T_\tau \times \mathbb{R}^{\dim(\tau)}) \times \left(\bigcup_{\gamma \in \mathcal{M}_\tau} \left(\theta(\text{int}(\gamma) \times [0, 1]) \times T^n / (\pi(\delta(\sigma_\gamma)) \times T_\tau) \right) \cup v \right).$$

Using the fact that θ is a topological embedding gives us

$$p^{-1}(\mathcal{U}) \cong (T_\tau \times \mathbb{R}^{\dim(\tau)}) \times \left([0, 1] \times \bigcup_{\gamma \in \mathcal{M}_\tau} \left(\text{int}(\gamma) \times T^n / (\pi(\delta(\sigma_\gamma)) \times T_\tau) \right) \cup v \right).$$

Thus, we have

$$(6) \quad p^{-1}(\mathcal{U}) \cong (T_\tau \times \mathbb{R}^{\dim(\tau)}) \times \mathcal{C} \left(\bigcup_{\gamma \in \mathcal{M}_\tau} \text{int}(\gamma) \times T^n / (\pi(\delta(\sigma_\gamma)) \times T_\tau) \right).$$

This homeomorphism gives us more than just the required *local* triviality. Recall that

$$X_{2j} - X_{2(j-1)} = p^{-1}(\mathcal{P}_j) - p^{-1}(\mathcal{P}_{j-1})$$

is simply the disjoint union of preimages of the interior of all j -dimensional faces of \mathcal{P} . Hence, we have

$$\begin{aligned} X_{2j} - X_{2(j-1)} &= \bigsqcup_{\substack{\tau \in \mathcal{P} \\ \dim(\tau)=j}} p^{-1}(\text{int}(\tau)) = \bigsqcup_{\substack{\tau \in \mathcal{P} \\ \dim(\tau)=j}} \text{int}(\tau) \times T^n / \pi(\delta(\tau)) \\ &\cong \bigsqcup_{\substack{\tau \in \mathcal{P} \\ \dim(\tau)=j}} \mathbb{R}^{\dim(\tau)} \times T_\tau. \end{aligned}$$

This means that $\mathbb{R}^{\dim(\tau)} \times T_\tau$ is a connected component of $X_{2j} - X_{2(j-1)}$. Thus, if we show that the given filtration endows $X_{\mathcal{P}}$ with a stratification then the link bundle is trivial. Now, consider the filtration of \mathcal{L}_τ that we introduced earlier

$$\mathcal{L} = \mathcal{L}_{2m+1} \supset \mathcal{L}_{2(m-1)+1} \supset \cdots \supset \mathcal{L}_1.$$

Let $x \in X_j - X_{j-1}$ and $x \in \text{int}(\tau)$. Thus, we can write $j = 2 \dim(\tau)$. Choose \mathcal{U} as described above. We want to investigate the intersection of $p^{-1}(\mathcal{U})$ with $X_{2 \dim(\tau)+i+1}$. Note that in the above filtration of toric varieties we have only even-dimensional¹ topological spaces X_j . Thus,

¹Note that X_{2j} can be identified with X_{2j+1} . This means that $X_{2j+1} - X_{2j} = \emptyset$ and thus the required conditions in Definition 10 are trivially fulfilled.

we can write $i + 1 = 2l$ with $l \in \mathbb{N}_{>0}$. Hence, we have

$$\begin{aligned}
& p^{-1}(\mathcal{U}) \cap X_{2(\dim(\tau)+l)} = \\
& \left(\bigcup_{\substack{\sigma \in \mathcal{P} \\ \dim(\sigma) \leq \dim(\tau)+l}} (\text{int}(\sigma) \times T^n / \pi(\delta(\sigma))) \right) \cap \\
& \left[(\mathbb{R}^{\dim(\tau)} \times T_\tau) \times \left(([0, 1] \times \bigcup_{\gamma \in \mathcal{M}_\tau} \text{int}(\gamma) \times T^n / (\pi(\delta(\sigma_\gamma)) \times T_\tau)) \cup v \right) \right] = \\
& (\mathbb{R}^{\dim(\tau)} \times T_\tau) \times \left[[0, 1] \times \left(\bigcup_{\substack{\sigma \in \mathcal{P} \\ \dim(\sigma) \leq \dim(\tau)+l}} \text{int}(\sigma) \times T^n / \pi(\delta(\sigma)) \right) \right. \\
& \left. \cap \left(\bigcup_{\gamma \in \mathcal{M}_\tau} \text{int}(\gamma) \times T^n / (\pi(\delta(\sigma_\gamma)) \times T_\tau) \right) \cup v \right] = \\
& (\mathbb{R}^{\dim(\tau)} \times T_\tau) \times \left[[0, 1] \times \left(\bigcup_{\substack{\gamma \in \mathcal{M}_\tau \\ \dim(\gamma) \leq l-1}} \text{int}(\gamma) \times T^n / (\pi(\delta(\sigma_\gamma)) \times T_\tau) \right) \cup v \right] = \\
& (\mathbb{R}^{\dim(\tau)} \times T_\tau) \times \mathcal{C}(\underbrace{(\mathcal{L}_\tau)_{2l-1}}_i).
\end{aligned}$$

Note that if $l = 0$ then we have

$$p^{-1}(\mathcal{U}) \cap X_{2\dim(\tau)} = (\mathbb{R}^{\dim(\tau)} \times T_\tau) \times v.$$

Consider that $|\mathcal{P}| - |\mathcal{P}^{n-1}|$ is the interior of $|\mathcal{P}|$ and it is dense in $|\mathcal{P}|$. This implies that $X_{\mathcal{P}} - X_{2(n-1)} \cong T^n \times \text{int}(\mathcal{P})$ is dense in $X_{\mathcal{P}}$. Now, if $\dim(\tau) = 0$ then $\mathcal{V} \cong \mathbb{R}^{\dim(|\mathcal{P}|)}$ and \mathcal{U} can be chosen as $\mathcal{C}(|\mathcal{M}_\tau|)$. This yields

$$p^{-1}(\mathcal{U}) \cong \mathcal{C}\left(\bigcup_{\gamma \in \mathcal{M}_\tau} \text{int}(\gamma) \times T^n / \pi(\delta(\sigma_\gamma))\right).$$

□

Remark 20. Note that $\mathcal{S}^1 \times \mathbb{R} \cong \mathbb{C}^*$. So $p^{-1}(\text{int}(\tau)) = (\mathbb{C}^*)^{\dim(\tau)}$ and specially

$$p^{-1}(\text{int}(\mathcal{P})) = (\mathbb{C}^*)^{\dim(|\mathcal{P}|)}.$$

There is an algebraic action of $(\mathbb{C}^*)^n$ on $X_{\mathcal{P}}$ with finitely many orbits. The preimage $p^{-1}(\text{int}(\mathcal{P}))$ is dense in $X_{\mathcal{P}}$, as mentioned earlier. One should also note that orbits are in one-to-one correspondence with the faces of \mathcal{P} and each stratum can be written as a disjoint union of finitely many orbits.

Remark 21. Given $\tau, \eta \in \mathcal{P}$ with $\dim(\tau) = \dim(\eta)$, \mathcal{L}_τ and \mathcal{L}_η are not necessarily even homotopy equivalent.

Example 22 ($n = 2$). Consider the complete fan introduced in Example 8. Let $\tau = \delta^{-1}(\tau_1)$ be the 1-dimensional dual face to τ_1 in \mathcal{P} . Consequently, $\mathcal{M}_\tau = \{\{0\}\}$ and $|\mathcal{M}_\tau| \cong *$. Note that $|\mathcal{M}_\tau| \subset \text{int}(\mathcal{P})$. Using the map defined in Expression 3 gives us

$$\begin{aligned}
\mathcal{M}_\tau \times T^2 &\longrightarrow \mathcal{L}_\tau \\
* \times T^2 &\longmapsto * \times T^2 / \mathcal{S}^1.
\end{aligned}$$

Hence, we get $\mathcal{L}_\tau \cong \mathcal{S}^1$. Now, let $\nu_{12} = \delta^{-1}(\sigma_{12})$ be the dual 0-dimensional face to σ_{12} in \mathcal{P} . The set $\mathcal{M}_{\nu_{12}}$ has 3 elements and $|\mathcal{M}_{\nu_{12}}| \cong \mathcal{I}$ with $\text{int}(\mathcal{M}_{\nu_{12}}) \subset \text{int}(\mathcal{P})$. We embed $\{0\} \in |\mathcal{M}_\tau|$ in

$|\mathcal{P}|$ such that $\{0\} \subset \text{int}(\delta^{-1}(\tau_1))$ and similarly $\{1\} \subset \text{int}(\delta^{-1}(\tau_2))$. Accordingly, we can describe $\mathcal{L}_{\nu_{12}}$ as below

$$\mathcal{L}_{\nu_{12}} \cong (\text{int}(\mathcal{I}) \times T^2) \cup (\{0\} \times T^2/\pi(\tau_1)) \cup (\{1\} \times T^2/\pi(\tau_2)).$$




FIGURE 5. Link of ν_{12} .

Remark 23. Note that $\mathcal{L}_{\nu_{12}} \not\cong \mathcal{S}^1 \times \mathcal{S}^2$. This can be easily deduced from the structure of σ_{12} or simply by comparing the homology groups computed later in this section. However, we can generalize our observation in the following form:

Let ν be a vertex of a 2-dimensional rational convex polytope \mathcal{P} with dual fan $\Sigma_{\mathcal{P}}$. Then ν has only 3 higher dimensional neighboring faces, namely two 1-dimensional faces τ and η and the 2-dimensional face, $\text{int}(\mathcal{P})$. Thus, we have

$$\mathcal{L}_{\nu_{12}} \cong (\text{int}(\mathcal{I}) \times \mathcal{T}^2) \cup (\{0\} \times \mathcal{T}^2/\pi(\delta(\eta))) \cup (\{1\} \times \mathcal{T}^2/\pi(\delta(\tau))).$$


and $\mathcal{L}_{\nu_{12}} \not\cong \mathcal{S}^1 \times \mathcal{S}^2$, which comes from the fact that we consider only complete proper fans.

Remark 24. Let \mathcal{P} be an n -dimensional rational polytope and τ an $(n-1)$ -dimensional face of \mathcal{P} . So $\mathcal{M}_{\tau} = \{\{0\}\}$, because $\text{int}(\mathcal{P})$ is the only higher dimensional neighboring face of τ . This implies that $\mathcal{L}_{\tau} \cong T^n/T^{n-1} \cong \mathcal{S}^1$. Later, we will use this observation and conclude that:

Let $X_{\mathcal{P}}$ be the toric variety associated to \mathcal{P} . Then $X_{\mathcal{P}}$ can not have a singular stratum with co-dimension 2.

Example 25 ($n = 3$). Consider the complete fan shown in Figure 6. We define the 1-dimensional cones as follows:


$$\begin{aligned} \tau_1 &= \{x(1, 0, 1) \mid x \in \mathbb{R}_{\geq 0}\}, \\ \tau_2 &= \{x(0, 1, 1) \mid x \in \mathbb{R}_{\geq 0}\}, \\ \tau_3 &= \{x(-1, 0, 1) \mid x \in \mathbb{R}_{\geq 0}\}, \\ \tau_4 &= \{x(0, -1, 1) \mid x \in \mathbb{R}_{\geq 0}\}, \\ \tau_5 &= \{x(0, 0, -1) \mid x \in \mathbb{R}_{\geq 0}\}. \end{aligned}$$

FIGURE 6. A complete fan Σ shown in \mathbb{R}^3 .

Let σ_{14} , illustrated in the above figure, be the 2-dimensional cone generated by the generators of τ_1 and τ_4 . We define the rest of the 2-dimensional cones similarly. At last, let ω_{1234} be the 3-dimensional cone which is generated by the generators of τ_1, τ_2, τ_3 , and τ_4 . The cones ω_{125} , ω_{235} , ω_{345} and ω_{145} are defined similarly. Consequently, we have

$$\Sigma = \{\{0\}, \tau_1, \tau_2, \tau_3, \tau_4, \tau_5, \sigma_{12}, \sigma_{23}, \sigma_{34}, \sigma_{14}, \omega_{125}, \omega_{235}, \omega_{345}, \omega_{145}, \omega_{1234}\}.$$

The polytope dual to Σ is then the pyramid shown in Figure 7.

FIGURE 7. The dual polytope of Σ illustrated in Figure 6.

As we discussed earlier, the link of a 2-dimensional face of \mathcal{P}_Σ is homeomorphic to \mathcal{S}^1 . Now, let $\eta_{12} = \delta^{-1}(\sigma_{12})$ be the 1-dimensional face of \mathcal{P} which is dual to σ_{12} . It is easy to see that $\mathcal{M}_{\eta_{12}} = \{\gamma_{\text{int}(\mathcal{P})}, \gamma_{\delta^{-1}(\tau_1)}, \gamma_{\delta^{-1}(\tau_2)}\}$. As in the 2-dimensional case we have $|\mathcal{M}_{\eta_{12}}| \cong \mathcal{I}$, where $\text{int}(\mathcal{M}_{\eta_{12}}) \subset \text{int}(\mathcal{P})$, and $\{0\} \subset \text{int}(\delta^{-1}(\tau_1))$ considered as a 0-dimensional face of \mathcal{I} . Similarly, we have $\{1\} \subset \text{int}(\delta^{-1}(\tau_2))$. This gives us

$$\mathcal{L}_{\eta_{12}} \cong (\text{int}(\mathcal{I}) \times T^3 / (T^3 / \pi(\sigma_{12}))) \cup (\{0\} \times T^3 / (T^3 / \pi(\sigma_{12}) \times \pi(\tau_1))) \cup (\{1\} \times T^3 / (T^3 / \pi(\sigma_{12}) \times \pi(\tau_2))).$$

Note that $T^3/(T^3/\pi(\sigma_{12})) \cong T^2$. Hence, we have

$$\mathcal{L}_{\eta_{12}} \cong (\text{int}(\mathcal{I}) \times T^2) \cup (\{0\} \times T^2/\pi(\tau_1)) \cup (\{1\} \times T^2/\pi(\tau_2)).$$

With the same argument as in the 2-dimensional case, we can show that $\mathcal{L}_{\eta_{12}} \not\cong \mathcal{S}^1 \times \mathcal{S}^2$. Forthwith, we want to describe a link of the point v at the apex of the pyramid. The set M_v is a 2-dimensional abstract polytope with four 1-dimensional faces and hence four 0-dimensional faces. Thus, we have

$$\mathcal{L}_v \cong (\text{int}(\mathcal{M}_v) \times T^3) \cup \bigcup_{\substack{\gamma_{\tau_i} \in \mathcal{M}_v \\ \dim(\gamma_{\tau_i})=1}} (\text{int}(\gamma_{\tau_i}) \times T^3/\pi(\delta(\tau_i))) \cup \bigcup_{\substack{\gamma_{\sigma_i} \in \mathcal{M}_v \\ \dim(\gamma_{\sigma_i})=0}} (\text{int}(\gamma_{\sigma_i}) \times T^3/\pi(\delta(\sigma_i))).$$

Recall that $T^3/\pi(\delta(\sigma_i)) \cong \mathcal{S}^1$ if $\dim(\sigma_i) = 1$. However, it is easy to show that we can not factor out any \mathcal{S}^1 in \mathcal{L}_v . The situation is displayed in Figure 8:




FIGURE 8. Link of the rationally singular point ν in X_Σ .

4. CW STRUCTURES ON TORIC VARIETIES

In this section, we endow toric varieties with CW structures. In his dissertation [19], Fischli has described a procedure that yields a CW structure of toric varieties. He carried out his method only for real 4-dimensional toric varieties. Our primary focus is on real 6-dimensional toric varieties. The path that we follow is slightly different from Fischli's, and one can use it also for real 6-dimensional compact toric varieties. The crucial idea is to ensure that each collapse $T^n/\pi(\tau) \rightarrow T^n/\pi(\sigma)$ for $\tau \prec \sigma$ with $\sigma, \tau \in \Sigma$, an n -dimensional complete rational fan, is cellular. We will see that in fact, the collapses are automatically cellular for 4-dimensional toric varieties. However, a slight modification is needed for 6-dimensional toric varieties. We will in addition give an idea of how this procedure can inductively be used for arbitrary dimensions. Again, we think of T^n as $\mathbb{R}^n/\mathbb{Z}^n$ and we continue to use the natural projection $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^n/\mathbb{Z}^n$. Let $\sigma \in \Sigma$ be a k -dimensional (rational) cone. There is, in addition, the orthogonal projection $\psi : \mathbb{R}^n \rightarrow \sigma^\perp$, where $\sigma^\perp = \{x \in \mathbb{R}^d \mid x \cdot y = 0 \ \forall y \in \sigma\}$. Hence, we can write $T^n/\pi(\sigma) \cong \sigma^\perp/\psi(\mathbb{Z}^n)$. Thus, choosing a CW structure on σ^\perp which is periodic with respect to $\psi(\mathbb{Z}^n)$ induces a finite CW structure on $T^n/\pi(\sigma)$. Closed n -cells homeomorphic to \mathcal{D}^n will be denoted by e^n .

4.1. Real 4-dimensional Toric Varieties. Let us start with real 4-dimensional toric varieties. We endow T^2 with the minimal CW structure with one 0-cell, two 1-cells and one 2-cell:

$$T^2 = e^0 \cup (e_{T_x}^1 \cup e_{T_y}^1) \cup_{f=xyx^{-1}y^{-1}} e_{T^2}^2$$

This structure is induced by the following CW structure on \mathbb{R}^2 :

On \mathbb{R} , each interval $[n, n+1]$ is considered as a 1-cell and each point $(n) \in \mathbb{R}$ is viewed as a 0-cell, where $n \in \mathbb{Z}$. The space \mathbb{R}^2 is then equipped with the product CW structure on $\mathbb{R} \times \mathbb{R}$. Now, let τ be a 1-dimensional cone in Σ which is generated by $\begin{pmatrix} n \\ m \end{pmatrix}$ where n and m are relatively

prime. Note that $\mathbb{R}^2 = \tau \oplus \tau^\perp$. Thus, we have then the following decomposition of $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$:

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = (-m) \begin{pmatrix} -m \\ m^2+n^2 \\ n \\ m^2+n^2 \end{pmatrix} + (n) \begin{pmatrix} n \\ m^2+n^2 \\ m \\ m^2+n^2 \end{pmatrix}.$$

Since $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ corresponds to the 1-cell $e_{T_y}^1$ of T^2 , the first coefficient, $-m$, of this decomposition will tell us the attaching degree of $\partial(e_{T_y}^1 \times e_{\mathcal{P}}^2)$ to cells $e_\tau^1 \times e_{\mathcal{P}}^1$. Now, consider the following CW structure on τ^\perp :

A 1-cell starts at the point $i \cdot \begin{pmatrix} -m \\ m^2+n^2 \\ n \\ m^2+n^2 \end{pmatrix}$ and ends at $(i+1) \cdot \begin{pmatrix} -m \\ m^2+n^2 \\ n \\ m^2+n^2 \end{pmatrix}$ for $i \in \mathbb{Z}$. Each point $i \cdot \begin{pmatrix} -m \\ m^2+n^2 \\ n \\ m^2+n^2 \end{pmatrix}$ is considered to be a 0-cell. This CW structure of τ^\perp induces a CW structure on $T^2/\pi(\tau)$. For each $\tau \in \Sigma$ equip $T^2/\pi(\tau)$ with the above CW structure, which has the form $T^2/\pi(\tau) = e_\tau^0 \cup e_\tau^1$. The CW structure on the polytope \mathcal{P} has been described in Section 2. (It is regular and consists of the faces of \mathcal{P} .) Consequently, we have the following cellular chain groups for the toric variety X_Σ :

$$\begin{aligned} \mathcal{C}_4(X) &= \mathbb{Q} \langle e_{T^2}^2 \times e_{\mathcal{P}}^2 \rangle \\ \mathcal{C}_3(X) &= \mathbb{Q} \langle e_{T^2}^1 \times e_{\mathcal{P}}^2 \rangle \oplus \mathbb{Q} \langle e_{T^2}^1 \times e_{\mathcal{P}}^2 \rangle \\ \mathcal{C}_2(X) &= \mathbb{Q} \langle e_{T^2}^0 \times e_{\mathcal{P}}^2 \rangle \oplus \bigoplus_{\substack{\tau_i \in \Sigma \\ \dim(\tau_i)=1}} \mathbb{Q} \langle e_{\tau_i}^1 \times e_{\mathcal{P}_i}^1 \rangle \\ \mathcal{C}_1(X) &= \bigoplus_{\substack{\tau_i \in \Sigma \\ \dim(\tau_i)=1}} \mathbb{Q} \langle e_{\tau_i}^0 \times e_{\mathcal{P}_i}^1 \rangle \\ \mathcal{C}_0(X) &= \bigoplus_{\substack{\sigma_i \in \Sigma \\ \dim(\sigma_i)=2}} \mathbb{Q} \langle e_{\sigma_i}^0 \times e_{\mathcal{P}_i}^0 \rangle. \end{aligned}$$

We keep in mind that $p^{-1}(\text{int}(\delta^{-1}(\tau_i))) \cong T^2/\pi(\tau_i) \times \text{int}(\delta^{-1}(\tau_i))$ and $\text{int}(\tau_i)$ represents the interior of a 1-cell, which we will denote by $e_{\mathcal{P}_i}^1$ in the above CW structure.

It remains to determine the boundary operators in the above CW structure. We can write $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ in the following form

$$\begin{pmatrix} 0 \\ 1 \end{pmatrix} = (n) \begin{pmatrix} -m \\ m^2+n^2 \\ n \\ m^2+n^2 \end{pmatrix} + (m) \begin{pmatrix} n \\ m^2+n^2 \\ m \\ m^2+n^2 \end{pmatrix}.$$

Since $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ corresponds to the 1-cell $e_{T_x}^1$ of T^2 , the first coefficient, n , of this decomposition will tell us the attaching degree of $\partial(e_{T_x}^1 \times e_{\mathcal{P}}^2)$ to cells $e_\tau^1 \times e_{\mathcal{P}}^1$. Let τ_i be an 1-dimensional cone in

Σ with $\begin{pmatrix} m_i \\ n_i \end{pmatrix}$ as the generator. In consequence, we have

$$\begin{aligned}\partial(e_{T^2}^1 \times e_{\mathcal{P}}^2) &= \sum_i n_i (e_{\tau_i}^1 \times e_{\mathcal{P}_i}^1) \\ \partial(e_{T^2}^1 \times e_{\mathcal{P}}^2) &= \sum_i -m_i (e_{\tau_i}^1 \times e_{\mathcal{P}_i}^1).\end{aligned}$$

At this point, we can compute the homology groups of the link of the point $x \in (X_\Sigma)_0$. Let σ_x be the dual cone to x in Σ . Let τ_1 and τ_2 be the 1-dimensional cones with $\tau_1, \tau_2 \prec \sigma$. Then, we have the following chain group for the link of x , \mathcal{L}_x .

$$\begin{aligned}\mathcal{C}_3(\mathcal{L}_x) &= \mathbb{Q} \langle e_{\mathcal{I}}^1 \times e_{T^2}^2 \rangle, \\ \mathcal{C}_2(\mathcal{L}_x) &= \mathbb{Q} \langle e_{\mathcal{I}}^1 \times e_{T^2}^1 \rangle \oplus \mathbb{Q} \langle e_{\mathcal{I}}^1 \times e_{T^2}^1 \rangle, \\ \mathcal{C}_1(\mathcal{L}_x) &= \mathbb{Q} \langle e_{\mathcal{I}}^1 \times e_{T^2}^0 \rangle \oplus \mathbb{Q} \langle e_{\mathcal{I}}^0 \times e_{S_1^0}^1 \rangle \oplus \mathbb{Q} \langle e_{\mathcal{I}}^0 \times e_{S_1^1}^1 \rangle, \\ \mathcal{C}_0(\mathcal{L}_x) &= \mathbb{Q} \langle e_{\mathcal{I}}^0 \times e_{S_1^0}^0 \rangle \oplus \mathbb{Q} \langle e_{\mathcal{I}}^0 \times e_{S_1^1}^0 \rangle.\end{aligned}$$

(Here the interval \mathcal{I} is the one defined in the Example 22.) We get the following boundary operators

$$(7) \quad \partial_3 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad \partial_2 = \begin{pmatrix} 0 & 0 \\ -m_1 & n_1 \\ -m_2 & n_2 \end{pmatrix}, \quad \partial_1 = \begin{pmatrix} +1 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix},$$

where $\begin{pmatrix} m_1 \\ n_1 \end{pmatrix}$ and $\begin{pmatrix} m_2 \\ n_2 \end{pmatrix}$ are the generators of τ_1 and τ_2 , respectively. Hence, we have

$$\begin{aligned}\text{rk}(H_3(\mathcal{L}_x; \mathbb{Q})) &= 1, \\ \text{rk}(H_2(\mathcal{L}_x; \mathbb{Q})) &= 0, \\ \text{rk}(H_1(\mathcal{L}_x; \mathbb{Q})) &= 0, \\ \text{rk}(H_0(\mathcal{L}_x; \mathbb{Q})) &= 1.\end{aligned}$$

As expected, \mathcal{L}_x is a rational homology sphere. But at this point, it is also worthwhile to study the homology groups of \mathcal{L}_x with integral coefficients. It is easy to see that $H_1(\mathcal{L}_x; \mathbb{Z}) = 0$ and only if $\det \begin{pmatrix} -m_1 & n_1 \\ -m_2 & n_2 \end{pmatrix} = \pm 1$. Thus, \mathcal{L}_x is an (integral) homology sphere if the previous condition holds. This means that X_σ is smooth if the generators of all $\sigma \in \Sigma$ with $\dim(\sigma) = 2$ satisfy the previous condition. However, there is also an algebraic description of the singularities of toric varieties. Cox, Little, and Schenck in [15, Theorem 1.3.12] show that a toric variety X_σ is smooth as a variety if and only if the minimal generators of $\sigma \subset \mathcal{N} \otimes \mathbb{R}$ form a part of an \mathbb{Z} -basis of the lattice \mathcal{N} for each $\sigma \in \Sigma$. Note that for real 4-dimensional toric varieties, this translates to the same condition that we obtained from our topological approach.

4.2. Real 6-dimensional Toric Varieties. In this section, we give a CW structure on real 6-dimensional toric varieties. As mentioned earlier, we need a slight refinement of the minimal CW structure of each T^2 here. The goal is to ensure that each collapse $T^3/\pi(\tau) \rightarrow T^3/\pi(\sigma)$, where $\tau \prec \sigma$ with $\sigma, \tau \in \Sigma$, is cellular. Here again, Σ is a complete fan.

At first, we consider the case of a 1-dimensional cone τ in Σ which is generated by $\begin{pmatrix} n \\ m \\ l \end{pmatrix}$,

where n, m and l are all nonzero and $\gcd(n, m, l) = 1$. Thus, we have $\tau = \text{span} \begin{pmatrix} n \\ m \\ l \end{pmatrix}$ and

$\tau^\perp = \text{span} \left(\begin{pmatrix} -m \\ n \\ 0 \end{pmatrix}, \begin{pmatrix} -l \\ 0 \\ n \end{pmatrix} \right)$. This gives us the following decomposition of $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = (-m) \begin{pmatrix} \frac{-m}{\Delta} \\ \frac{n}{\Delta} \\ 0 \end{pmatrix} + (-l) \begin{pmatrix} \frac{-l}{\Delta} \\ 0 \\ \frac{n}{\Delta} \end{pmatrix} + (n) \begin{pmatrix} \frac{n}{\Delta} \\ \frac{m}{\Delta} \\ \frac{l}{\Delta} \end{pmatrix},$$

where $\Delta = l^2 + n^2 + m^2$. We can write the above decomposition as

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = (lm) \begin{pmatrix} \frac{m^2+l^2}{\Delta} \frac{1}{\Delta} \\ \frac{lm}{\Delta} \frac{1}{\Delta} \\ -\frac{nl}{\Delta} \frac{1}{\Delta} \end{pmatrix} + (n) \begin{pmatrix} \frac{n}{\Delta} \\ \frac{m}{\Delta} \\ \frac{l}{\Delta} \end{pmatrix}.$$

Similarly, we have

$$\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = (nl) \begin{pmatrix} \frac{-nm}{\Delta} \frac{1}{\Delta} \\ \frac{n^2+l^2}{\Delta} \frac{1}{\Delta} \\ -\frac{ml}{\Delta} \frac{1}{\Delta} \end{pmatrix} + (m) \begin{pmatrix} \frac{n}{\Delta} \\ \frac{m}{\Delta} \\ \frac{l}{\Delta} \end{pmatrix}$$

$$\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = (nm) \begin{pmatrix} \frac{-nl}{\Delta} \frac{1}{\Delta} \\ -\frac{ml}{\Delta} \frac{1}{\Delta} \\ \frac{n^2+m^2}{\Delta} \frac{1}{\Delta} \end{pmatrix} + (l) \begin{pmatrix} \frac{n}{\Delta} \\ \frac{m}{\Delta} \\ \frac{l}{\Delta} \end{pmatrix}.$$

With these in hand, we endow τ^\perp with the following CW structure. The points $i \begin{pmatrix} \frac{m^2+l^2}{\Delta} \frac{1}{\Delta} \\ \frac{lm}{\Delta} \frac{1}{\Delta} \\ -\frac{nl}{\Delta} \frac{1}{\Delta} \end{pmatrix}$,

$i \begin{pmatrix} \frac{-nm}{\Delta} \frac{1}{\Delta} \\ \frac{n^2+l^2}{\Delta} \frac{1}{\Delta} \\ -\frac{ml}{\Delta} \frac{1}{\Delta} \end{pmatrix}$, and $i \begin{pmatrix} \frac{-nl}{\Delta} \frac{1}{\Delta} \\ -\frac{ml}{\Delta} \frac{1}{\Delta} \\ \frac{n^2+m^2}{\Delta} \frac{1}{\Delta} \end{pmatrix}$ are considered to be 0-cells where $i \in \mathbb{Z}$. A 1-cell starts at

$i \begin{pmatrix} \frac{m^2+l^2}{\Delta} \frac{1}{\Delta} \\ \frac{lm}{\Delta} \frac{1}{\Delta} \\ -\frac{nl}{\Delta} \frac{1}{\Delta} \end{pmatrix}$ and ends at $(i+1) \begin{pmatrix} \frac{m^2+l^2}{\Delta} \frac{1}{\Delta} \\ \frac{lm}{\Delta} \frac{1}{\Delta} \\ -\frac{nl}{\Delta} \frac{1}{\Delta} \end{pmatrix}$ (similarly for $\begin{pmatrix} \frac{-nm}{\Delta} \frac{1}{\Delta} \\ \frac{n^2+l^2}{\Delta} \frac{1}{\Delta} \\ -\frac{ml}{\Delta} \frac{1}{\Delta} \end{pmatrix}$ and $\begin{pmatrix} \frac{-nl}{\Delta} \frac{1}{\Delta} \\ -\frac{ml}{\Delta} \frac{1}{\Delta} \\ \frac{n^2+m^2}{\Delta} \frac{1}{\Delta} \end{pmatrix}$).

Note that

$$\begin{pmatrix} \frac{m^2+l^2}{\Delta} \frac{1}{\Delta} \\ \frac{lm}{\Delta} \frac{1}{\Delta} \\ -\frac{nl}{\Delta} \frac{1}{\Delta} \end{pmatrix} + \begin{pmatrix} \frac{-nm}{\Delta} \frac{1}{\Delta} \\ \frac{n^2+l^2}{\Delta} \frac{1}{\Delta} \\ -\frac{ml}{\Delta} \frac{1}{\Delta} \end{pmatrix} + \begin{pmatrix} \frac{-nl}{\Delta} \frac{1}{\Delta} \\ -\frac{ml}{\Delta} \frac{1}{\Delta} \\ \frac{n^2+m^2}{\Delta} \frac{1}{\Delta} \end{pmatrix} = 0.$$

The above relation ensures that we can form two 2-simplices with the above vectors. We consider these 2-simplices as 2-cells of τ^\perp . From the construction, it is clear that the previous CW structure is periodic with respect to $\psi(\mathbb{Z}^3)$, where ψ denotes the natural projection from \mathbb{R}^3 onto τ^\perp . The induced CW structure on $T^3/\pi(\tau) \cong \tau^\perp/\psi(\mathbb{Z}^3) \cong T^2$ can be described as follows.

$$(8) \quad T^2 = (e_{T^2}^2 \cup e_{T^2}^2) \cup (e_{T^2}^1 \cup e_{T^2}^1 \cup e_{T^2}^1) \cup e_{T^2}^0.$$

With an appropriate orientation we have the following boundary operators.

$$\partial_2^{T^2} = \begin{pmatrix} -1 & 1 \\ -1 & 1 \\ -1 & 1 \end{pmatrix}, \quad \partial_1^{T^2} = (0 \ 0 \ 0).$$

Schematically, we have




FIGURE 9. CW structure on T^2 with two 2-Cells. Note that we identify the opposite boundary edges as in the minimal CW structure on T^2 .

There is yet another case that we need to consider and study, namely when one or two of n, m, l are zero. Without loss of generality assume that $\tau = \text{span} \begin{pmatrix} n \\ m \\ 0 \end{pmatrix}$. This yields

$$\tau^\perp = \text{span} \left(\begin{pmatrix} -m \\ n \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right).$$

Applying the previous construction induces the following CW structure on $T^3/\pi(\tau) \cong T^2$.

$$T^2 = e_{T^2}^2 \cup (e_{T^2}^1 \cup e_{T^2}^1) \cup e_{T^2}^0,$$

which is the minimal CW structure on T^2 with the well-known vanishing boundary operators. From the previous section, we know that collapses of the form $T^2 \rightarrow \mathcal{S}^1$ are automatically cellular. Thus to ensure cellularity, it is enough to equip each $T^3/\pi(\tau) \cong T^2$ with the appropriate CW structure. To determine the boundary operators of X_Σ associated with this CW structure, one could also go further and investigate the collapses $T^3/\pi(\tau) \rightarrow T^3/\pi(\sigma)$, such that $\tau \prec \sigma$, $\dim(\tau) = 1$, $\dim(\sigma) = 2$ and $\tau, \sigma \in \Sigma$. Since we are primarily interested in rational homology, it will turn out that we do not need these collapsing data explicitly for our purposes.

For the sake of simplicity, we introduce the following notation that we will use for some of our matrix representations. Let \mathbf{A} be an $lp \times kh$ matrix where $l, p, k, h \in \mathbb{N}$. For our purposes, it is sometimes practical to study only a specific part of a matrix. Let us now consider \mathbf{A} as an $p \times h$ matrix where each element (or block) of \mathbf{A} is an $l \times k$ matrix. From the context it will be clear how one can obtain the rest of the matrix from an arbitrary block. Then we represent \mathbf{A} as

$$\mathbf{A} = \left(\begin{array}{c} \overbrace{\left\{ \right\}}^k \\ \underbrace{\left\{ \right\}}_l \end{array} \right)_{p \times h}.$$

At last, we aim to compute the homology groups of an arbitrary real 6-dimensional toric variety.

4.2.1. *The Homology of real 6-dimensional Toric Varieties.* Let Σ be a complete fan and \mathcal{P} the associated dual polytope to Σ . Let f_1, f_2 and f_3 denote the number of 1-dimensional, 2-dimensional and 3-dimensional cones of Σ , respectively. With the above considerations, we can endow $X_{\mathcal{P}}$, the toric variety associated to \mathcal{P} , with the following cellular chain groups:

$$\begin{aligned}
\mathcal{C}_6(X_{\mathcal{P}}) &= \mathbb{Q} \langle e_{T^3}^3 \times e_{\mathcal{P}}^3 \rangle \\
\mathcal{C}_5(X_{\mathcal{P}}) &= \bigoplus_{i=1}^3 \mathbb{Q} \langle e_{T_i^2}^2 \times e_{\mathcal{P}}^3 \rangle \\
\mathcal{C}_4(X_{\mathcal{P}}) &= \bigoplus_{i=1}^3 \mathbb{Q} \langle e_{T_i^2}^1 \times e_{\mathcal{P}}^3 \rangle \bigoplus_{i=1}^{\gamma} (\mathbb{Q} \langle e_{(T_{\gamma_i}^2)_1}^2 \times e_{\mathcal{P}_{\gamma_i}}^2 \rangle \oplus \mathbb{Q} \langle e_{(T_{\gamma_i}^2)_2}^2 \times e_{\mathcal{P}_{\gamma_i}}^2 \rangle) \\
&\quad \bigoplus_{j=1}^{\omega} \mathbb{Q} \langle e_{T_{\omega_j}^2}^2 \times e_{\mathcal{P}_{\omega_j}}^2 \rangle \\
\mathcal{C}_3(X_{\mathcal{P}}) &= \mathbb{Q} \langle e_{T^3}^0 \times e_{\mathcal{P}}^3 \rangle \bigoplus_{i=1}^{\gamma} \bigoplus_{l=1}^3 \mathbb{Q} \langle e_{(T_{\gamma_i}^2)_l}^1 \times e_{\mathcal{P}_{\gamma_i}}^2 \rangle \bigoplus_{j=1}^{\omega} \bigoplus_{l=1}^2 \mathbb{Q} \langle e_{(T_{\omega_j}^2)_l}^1 \times e_{\mathcal{P}_{\omega_j}}^2 \rangle \\
\mathcal{C}_2(X_{\mathcal{P}}) &= \bigoplus_{i=1}^{\gamma} \mathbb{Q} \langle e_{(T_{\gamma_i}^2)}^0 \times e_{\mathcal{P}_{\gamma_i}}^2 \rangle \bigoplus_{j=1}^{\omega} \mathbb{Q} \langle e_{(T_{\omega_j}^2)}^0 \times e_{\mathcal{P}_{\omega_j}}^1 \rangle \bigoplus_{l=1}^{f_2} \mathbb{Q} \langle e_{S_l^1}^1 \times e_{\mathcal{P}_l}^1 \rangle \\
\mathcal{C}_1(X_{\mathcal{P}}) &= \bigoplus_{i=0}^{f_1} \mathbb{Q} \langle e_{S_i^1}^0 \times e_{\mathcal{P}_i}^1 \rangle \\
\mathcal{C}_0(X_{\mathcal{P}}) &= \bigoplus_{i=0}^{f_3} \mathbb{Q} \langle e_{\mathcal{P}_i}^0 \rangle,
\end{aligned}$$

where γ and ω are the number of 2-dimensional tori with three and two 1-cells, respectively. The cell $e_{\gamma_i}^2$ is the 2-cell of \mathcal{P} , endowed with the regular CW structure, which is attached to $T_{\gamma_i}^2$, where $i = 1, \dots, \gamma$, and similarly for $e_{\mathcal{P}_{\omega_j}}^2$. Hence, we have $\gamma + \omega = f_1$.

Remark 26. *In the following discussion, without loss of generality and for the sake of simplicity, we only consider $\begin{pmatrix} n_{\omega_j} \\ m_{\omega_j} \\ 0 \end{pmatrix}$ as generators of 1-dimensional faces in Σ with at least one zero-entry.*

Considering

$$\begin{pmatrix} 0 \\ m_{\omega_j} \\ l_{\omega_j} \end{pmatrix}, \begin{pmatrix} n_{\omega_j} \\ 0 \\ l_{\omega_j} \end{pmatrix}, \begin{pmatrix} n_{\omega_j} \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ m_{\omega_j} \\ 0 \end{pmatrix}, \text{ and } \begin{pmatrix} 0 \\ 0 \\ l_{\omega_j} \end{pmatrix}$$

will merely change the position of zero elements in the rows labeled by ω in the boundary operators ∂_5 and ∂_4 . The computation goes along the same lines.

Let $\begin{pmatrix} n_{\gamma_i} \\ m_{\gamma_i} \\ l_{\gamma_i} \end{pmatrix}$ and $\begin{pmatrix} n_{\omega_i} \\ m_{\omega_i} \\ 0 \end{pmatrix}$, with $i = 1, \dots, \gamma$ and $j = 1, \dots, \omega$, be the generator of 1-dimensional cones in Σ dual to $e_{\mathcal{P}_{\gamma_i}}^2$ and $e_{\mathcal{P}_{\omega_j}}^2$, respectively, where we consider the 2-cells as 2-dimensional faces of \mathcal{P} . Consider the 2-cell, $e_{T_3}^2$, in $T^3 \cong \mathbb{R}^3/\mathbb{Z}^3$ with $e_{T_1}^1$ and $e_{T_2}^1$ in its topological boundary, where $e_{T_1}^1$ is the 1-cell of T^3 induced by $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ under the map $\mathbb{R}^3 \rightarrow \mathbb{R}^3/\mathbb{Z}^3$ and $e_{T_2}^1$ induced by $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$.

Under the projection $\mathbb{R}^3 \rightarrow \text{span}\left(\begin{pmatrix} n_{\gamma_i} \frac{1}{\Delta} \\ m_{\gamma_i} \frac{1}{\Delta} \\ l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix}\right)^\perp$, we map the vectors $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ to $\begin{pmatrix} (m_{\gamma_i}^2 + l_{\gamma_i}^2) \frac{1}{\Delta} \\ -m_{\gamma_i} n_{\gamma_i} \frac{1}{\Delta} \\ -n_{\gamma_i} l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix}$ and $\begin{pmatrix} -n_{\gamma_i} m_{\gamma_i} \frac{1}{\Delta} \\ (n_{\gamma_i}^2 + l_{\gamma_i}^2) \frac{1}{\Delta} \\ -m_{\gamma_i} l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix}$, respectively. Hence, the area enclosed by the vectors $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ maps to the 2-cell enclosed by $\begin{pmatrix} (m_{\gamma_i}^2 + l_{\gamma_i}^2) \frac{1}{\Delta} \\ -m_{\gamma_i} n_{\gamma_i} \frac{1}{\Delta} \\ -n_{\gamma_i} l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix}$ and $\begin{pmatrix} -n_{\gamma_i} m_{\gamma_i} \frac{1}{\Delta} \\ (n_{\gamma_i}^2 + l_{\gamma_i}^2) \frac{1}{\Delta} \\ -m_{\gamma_i} l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix}$ in $\text{span}\left(\left(\begin{pmatrix} n_{\gamma_i} \frac{1}{\Delta} \\ m_{\gamma_i} \frac{1}{\Delta} \\ l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix}\right)^\perp\right)$. We have

$$\left\| \begin{pmatrix} (m_{\gamma_i}^2 + l_{\gamma_i}^2) \frac{1}{\Delta} \\ -m_{\gamma_i} n_{\gamma_i} \frac{1}{\Delta} \\ -n_{\gamma_i} l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix} \times \begin{pmatrix} -n_{\gamma_i} m_{\gamma_i} \frac{1}{\Delta} \\ (n_{\gamma_i}^2 + l_{\gamma_i}^2) \frac{1}{\Delta} \\ -m_{\gamma_i} l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix} \right\| = \frac{l_{\gamma_i}}{\sqrt{\Delta}}.$$

Note that the area enclosed by the two generators in $\text{span}\left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}\right)$ is

$$\left\| \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \times \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right\| = 1.$$

However, the area enclosed by the generators of a 2-cell in $\text{span}\left(\left(\begin{pmatrix} n_{\gamma_i} \frac{1}{\Delta} \\ m_{\gamma_i} \frac{1}{\Delta} \\ l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix}\right)^\perp\right)$, since n_{γ_i} ,

m_{γ_i} , and l_{γ_i} are coprime, is $\left\| \begin{pmatrix} n_{\gamma_i} \frac{1}{\Delta} \\ m_{\gamma_i} \frac{1}{\Delta} \\ l_{\gamma_i} \frac{1}{\Delta} \end{pmatrix} \right\| = \frac{1}{\sqrt{\Delta}}$. It follows that the degree of the map from

the boundary of $e_{T_3}^2 \times e_{\mathcal{P}}^3$ to the boundaries of $e_{(T_{\gamma_i}^2)_1}^2 \times e_{\mathcal{P}_{\gamma_i}}^2$ and $e_{(T_{\gamma_i}^2)_2}^2 \times e_{\mathcal{P}_{\gamma_i}}^2$ is $\frac{l_{\gamma_i}/\sqrt{\Delta}}{1/\sqrt{\Delta}} = l_{\gamma_i}$. Note that $\omega + \gamma \geq 4$. Hence, we get the following boundary operators. If the generator has at least one zero entry, the treatment goes also along the same lines.

$$\partial_6 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\partial_5 = \begin{array}{c} e_{T_1^3}^1 \times e_{\mathcal{P}}^3 \\ e_{T_2^3}^1 \times e_{\mathcal{P}}^3 \\ e_{T_3^3}^1 \times e_{\mathcal{P}}^3 \\ e_{(T_{\gamma_i}^2)_1}^2 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{(T_{\gamma_i}^2)_2}^2 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{T_{\omega_j}^2}^2 \times e_{\mathcal{P}_{\omega_j}}^2 \omega \end{array} \begin{bmatrix} e_{T_1^3}^2 \times e_{\mathcal{P}}^3 & e_{T_2^3}^2 \times e_{\mathcal{P}}^3 & e_{T_3^3}^2 \times e_{\mathcal{P}}^3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ n_{\gamma_i} & m_{\gamma_i} & l_{\gamma_i} \\ n_{\gamma_i} & m_{\gamma_i} & l_{\gamma_i} \\ n_{\omega_j} & m_{\omega_j} & 0 \end{bmatrix}$$

$$\partial_4 = \begin{array}{c} e_{T_3}^0 \times e_{\mathcal{P}}^3 \\ e_{(T_{\gamma_i}^2)_1}^1 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{(T_{\gamma_i}^2)_2}^1 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{(T_{\gamma_i}^2)_3}^1 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{(T_{\omega_j}^2)_1}^1 \times e_{\mathcal{P}_{\omega_j}}^2 \omega \\ e_{(T_{\omega_j}^2)_2}^1 \times e_{\mathcal{P}_{\omega_j}}^2 \omega \end{array} \begin{bmatrix} e_{T_1^3}^1 \times e_{\mathcal{P}}^3 & e_{T_2^3}^1 \times e_{\mathcal{P}}^3 & e_{T_3^3}^1 \times e_{\mathcal{P}}^3 & e_{(T_{\gamma_i}^2)_1}^2 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma & e_{(T_{\gamma_i}^2)_2}^2 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma & e_{T_{\omega_j}^2}^2 \times e_{\mathcal{P}_{\omega_j}}^2 \omega \\ 0 & 0 & 0 & \overbrace{0}^{\gamma} & \overbrace{0}^{\gamma} & \overbrace{0}^{\omega} \\ m_{\gamma_i} l_{\gamma_i} & 0 & 0 & 1 & -1 & 0 \\ 0 & n_{\gamma_i} l_{\gamma_i} & 0 & 1 & -1 & 0 \\ 0 & 0 & n_{\gamma_i} m_{\gamma_i} & 1 & -1 & 0 \\ -m_{\omega_j} & 0 & 0 & 0 & 0 & 0 \\ 0 & n_{\omega_j} & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\partial_3 = \begin{array}{c} e_{T_1^3}^0 \times e_{\mathcal{P}}^3 \\ e_{(T_{\gamma_i}^2)_1}^0 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{(T_{\gamma_i}^2)_2}^0 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{(T_{\gamma_i}^2)_3}^0 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{(T_{\omega_j}^2)_1}^0 \times e_{\mathcal{P}_{\omega_j}}^2 \omega \\ e_{(T_{\omega_j}^2)_2}^0 \times e_{\mathcal{P}_{\omega_j}}^2 \omega \\ e_{S_1^1}^1 \times e_{\mathcal{P}_l}^1 f_2 \end{array} \begin{bmatrix} e_{T_1^3}^0 \times e_{\mathcal{P}}^3 & e_{(T_{\gamma_i}^2)_1}^1 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma & e_{(T_{\gamma_i}^2)_2}^1 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma & e_{(T_{\gamma_i}^2)_3}^1 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma & e_{(T_{\omega_j}^2)_1}^1 \times e_{\mathcal{P}_{\omega_j}}^2 \omega & e_{(T_{\omega_j}^2)_2}^1 \times e_{\mathcal{P}_{\omega_j}}^2 \omega \\ 1 & \overbrace{0}^{\gamma} & \overbrace{0}^{\gamma} & \overbrace{0}^{\gamma} & \overbrace{0}^{\omega} & \overbrace{0}^{\omega} \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & C_1 & C_2 & C_3 & C_4 & C_5 \end{bmatrix}$$

$$\partial_2 = \begin{array}{c} e_{T_{\gamma_i}^2}^0 \times e_{\mathcal{P}_{\gamma_i}}^2 \gamma \\ e_{T_{\omega_j}^2}^0 \times e_{\mathcal{P}_{\omega_j}}^2 \omega \\ e_{S_1^1}^1 \times e_{\mathcal{P}_l}^1 f_2 \end{array} \left\{ \begin{bmatrix} \overbrace{P_1}^{\gamma} & \overbrace{P_2}^{\omega} & \overbrace{0}^{f_2} \end{bmatrix} \right.$$

where C_i , $i = 1, \dots, 5$ are collapsing data that we do not need explicitly. The values in the blocks P_i for $i = 1, 2$ are purely determined by the regular CW structure on \mathcal{P} . Then, one can easily deduce that

$$\begin{aligned} \text{rk}(H_6(X_{\mathcal{P}}; \mathbb{Q})) &= 1, \\ \text{rk}(H_5(X_{\mathcal{P}}; \mathbb{Q})) &= 0. \end{aligned}$$

Moreover, we have

$$\begin{aligned}
\mathrm{rk}(\mathrm{Im}(\partial_5)) &= 3, \\
\mathrm{rk}(\ker(\partial_4)) &= \gamma + \omega, \\
\mathrm{rk}(\mathrm{Im}(\partial_4)) &= \gamma + 3, \\
\mathrm{rk}(\ker(\partial_2)) &= \mathrm{rk}(\ker(\partial_2^{\mathcal{P}})) + f_2, \\
\mathrm{rk}(\mathrm{Im}(\partial_2)) &= \mathrm{rk}(\mathrm{Im}(\partial_2^{\mathcal{P}})) \\
\partial_1 &= \partial_1^{\mathcal{P}}
\end{aligned}$$

where $\partial_i^{\mathcal{P}}$ denote the boundary operators of $\mathcal{P} \cong \mathcal{D}^3$. Thus, we have

$$\mathrm{rk}(H_4(X_{\mathcal{P}}; \mathbb{Q})) = f_1 - 3.$$

Define b' by the equation

$$(9) \quad \mathrm{rk}(\mathrm{Im}(\partial_3)) = f_2 - b' + \mathrm{rk}(\mathrm{Im}(\partial_3^{\mathcal{P}}))$$

and b by the equation

$$(10) \quad b' = f_1 - 3 - b.$$

In other words, b' is the number of linearly dependent rows of ∂_3 . Straightforward calculation then shows that

$$\begin{aligned}
\mathrm{rk}(\mathrm{Im}(\partial_3)) &= -(f_1 - f_2 - b - 3) + \mathrm{rk}(\mathrm{Im}(\partial_3^{\mathcal{P}})), \\
\mathrm{rk}(\ker(\partial_3)) &= (f_1 - f_2 - b - 3) + (3\gamma + 2\omega).
\end{aligned}$$

This implies that

$$\mathrm{rk}(H_3(X_{\mathcal{P}}; \mathbb{Q})) = 3f_1 - f_2 - b - 6.$$

and

$$\mathrm{rk}(H_2(X_{\mathcal{P}}; \mathbb{Q})) = f_1 - 3 - b.$$

If $X_{\mathcal{P}}$ is non-singular then Poincaré duality implies $b = 0$ and hence $b' = f_1 - 3$. The previous considerations lead to the following proposition.

Proposition 27. *Let $X_{\mathcal{P}}$ be a 6-dimensional toric variety associated to a complete rational fan Σ , which is dual to the polytope \mathcal{P} . Let f_2 and f_1 be the number of 2- and 1-dimensional cones in Σ , respectively. Then, we have*

$$\begin{aligned}
\mathrm{rk}(H_6(X_{\mathcal{P}}; \mathbb{Q})) &= 1, \\
\mathrm{rk}(H_5(X_{\mathcal{P}}; \mathbb{Q})) &= 0, \\
\mathrm{rk}(H_4(X_{\mathcal{P}}; \mathbb{Q})) &= f_1 - 3, \\
\mathrm{rk}(H_3(X_{\mathcal{P}}; \mathbb{Q})) &= 3f_1 - f_2 - b - 6, \\
\mathrm{rk}(H_2(X_{\mathcal{P}}; \mathbb{Q})) &= f_1 - 3 - b, \\
\mathrm{rk}(H_1(X_{\mathcal{P}}; \mathbb{Q})) &= 0, \\
\mathrm{rk}(H_0(X_{\mathcal{P}}; \mathbb{Q})) &= 1,
\end{aligned}$$

where the parameter b is determined by Equations 9 and 10.

Remark 28. *In the next section, we give a geometric description of b , at least for some special cases. There is yet another approach by McConnell in [27], which computes the homology groups of 6-dimensional toric varieties, using spectral sequences.*

Remark 29. *As observed by McConnell in [27], b is not combinatorially invariant, i.e. b is not merely determined by the number of cones.*

4.3. Singularities of 6-dimensional Toric Varieties. Let Σ be a complete fan in \mathbb{R}^3 and \mathcal{P} be the associated dual polytope. As mentioned in Remark 24, the link of a point $x \in X_4 - X_2$ is simply homeomorphic to \mathcal{S}^1 . For $x \in X_2 - X_0$, we employ the construction of links introduced earlier. Let $\tau \in \Sigma$ such that $p(x) \in \text{int}(\delta^{-1}(\tau))$, where p is the natural projection $p : X_\Sigma \rightarrow |\mathcal{P}|$. Then, we have $\mathcal{S}_\tau = \{\sigma_1, \sigma_2, \text{int}(|\mathcal{P}|)\}$, where σ_1 and σ_2 are the two 2-dimensional neighboring faces of τ . Hence as in Example 22, we have $\mathcal{M}_\tau \cong \mathcal{I}$. However, in contrast to Example 22, here we have $T^3/\pi(\delta(\tau)) \cong \mathcal{S}^1$. But in the end, \mathcal{L}_τ , the link of x , has the same CW structure as in Example 22,

$$\mathcal{L}_\tau \cong \left(\text{int}(\mathcal{I}) \times T^3 / (T^3 / \pi(\delta(\tau))) \right) \bigcup \left(\{0\} \times (T^3 / \delta(\pi(\sigma_1))) / (T^3 / \pi(\delta(\tau))) \right) \bigcup \left(\{1\} \times (T^3 / \delta(\pi(\sigma_2))) / (T^3 / \pi(\delta(\tau))) \right).$$

Note that with the same argument as in Remark 23, we conclude that $\mathcal{L} \not\cong \mathcal{S}^1 \times \mathcal{S}^2$. The computational method goes along the same line as in Section 4.1 and yields

$$(11) \quad \begin{aligned} \text{rk}(H_3(\mathcal{L}_\tau; \mathbb{Q})) &= 1, \\ \text{rk}(H_2(\mathcal{L}_\tau; \mathbb{Q})) &= 0, \\ \text{rk}(H_1(\mathcal{L}_\tau; \mathbb{Q})) &= 0, \\ \text{rk}(H_0(\mathcal{L}_\tau; \mathbb{Q})) &= 1. \end{aligned}$$

Remark 30. *Studying the homology groups of \mathcal{L}_τ with integral coefficients yields the same result for the smoothness of a 2-dimensional stratum in X_Σ as in the previous case. This means that if the two generators of $\delta(\tau)$ in $\Sigma = \mathcal{N} \otimes \mathbb{R}$ do not form a part of a \mathbb{Z} -basis of the lattice \mathcal{N} then the stratum is singular in the sense of Definition 13.*

In Example 25, we constructed the link of a connected component of X_0 (a point) in a specific real 6-dimensional toric variety. Here, we generalize this Example to an arbitrary real 6-dimensional compact toric variety.

Let $x \in X_0$. We consider x as a 0-dimensional face of \mathcal{P} . The set \mathcal{S}_x consists of $\text{int}(\mathcal{P})$ and all 1- and 2-dimensional faces of \mathcal{P} , τ , with $x \prec \tau$. Following the introduced construction of links, $|\mathcal{M}_x|$ is a 2-dimensional polytope with f_{x_1} vertices and f_{x_2} 1-dimensional faces, where f_{x_1} and f_{x_2} denote the number of 1- and 2-dimensional neighboring faces of x in Σ , respectively. Since $|\mathcal{M}_x| \cong \mathcal{D}^2$, its boundary is homeomorphic to a circle. As the Euler characteristic of a circle vanishes, $f_{x_1} = f_{x_2}$. Hence, we have the following relations.

$$\begin{aligned} p_{\mathcal{L}_x}^{-1}(\text{int}(\mathcal{M}_x)) &= T^3 \times \text{int}(\mathcal{M}_x) \\ p_{\mathcal{L}_x}^{-1}(\text{int}(\tau)) &= T^2 \times \text{int}(\tau) \quad \text{for } \tau \in \mathcal{M}_x \text{ with } \dim(\tau) = 1 \\ p_{\mathcal{L}_x}^{-1}(\nu) &= \mathcal{S}^1 \times \nu \quad \text{for } \nu \in \mathcal{M}_x \text{ with } \dim(\nu) = 0. \end{aligned}$$

At this point, we can endow \mathcal{L}_x with a CW structure and compute the corresponding homology groups. As before, we equip T^3 with the minimal CW structure and each T^2 with an appropriate CW structure such that each collapse map becomes cellular as we discussed earlier. Each 1-dimensional face of \mathcal{M}_x , τ_i , is associated to a 2-dimensional face of \mathcal{P} , which is dual to a 1-dimensional cone in Σ such that the dual cones lie in \mathcal{S}_x . Let a and b be the numbers of such 1-dimensional cones in Σ , whose generators have no zero entry and at least one zero-entry,

respectively. Take into consideration that $a + b \geq 3$. Let $\begin{pmatrix} n_{a_i} \\ m_{a_i} \\ l_{a_i} \end{pmatrix}$ and $\begin{pmatrix} n_{b_j} \\ m_{b_j} \\ 0 \end{pmatrix}$ with $i = 1, \dots, a$

and $j = 1, \dots, b$ be the generators of these 1-dimensional cones in Σ where n_{a_i}, m_{a_i} and l_{a_i} are non-zero and at least one of the entries n_{b_j} or m_{b_j} is non-zero.

Accordingly, we get the following chain groups for \mathcal{L}_x , where we use the regular cellular chain groups on \mathcal{M}_x :

$$\begin{aligned}
\mathcal{C}_5(\mathcal{L}_x) &= \mathbb{Q} \langle e_{T^3}^3 \times e_{\mathcal{M}_x}^2 \rangle \\
\mathcal{C}_4(\mathcal{L}_x) &= \bigoplus_{i=1}^3 \mathbb{Q} \langle e_{T_i^3}^2 \times e_{\mathcal{M}_x}^2 \rangle \\
\mathcal{C}_3(\mathcal{L}_x) &= \bigoplus_{i=1}^3 \mathbb{Q} \langle e_{T_i^3}^1 \times e_{\mathcal{M}_x}^2 \rangle \bigoplus_{i=1}^a (\mathbb{Q} \langle e_{(T_{a_i}^2)_1}^2 \times e_{(\mathcal{M}_{x_a})_i}^1 \rangle \oplus \mathbb{Q} \langle e_{(T_{a_i}^2)_2}^2 \times e_{(\mathcal{M}_{x_a})_i}^1 \rangle) \\
&\quad \bigoplus_{j=1}^b \mathbb{Q} \langle e_{(T_{b_j}^2)}^2 \times e_{(\mathcal{M}_{x_b})_j}^1 \rangle \\
\mathcal{C}_2(\mathcal{L}_x) &= \mathbb{Q} \langle e_{T^3}^0 \times e_{\mathcal{M}_x}^2 \rangle \bigoplus_{i=1}^a \bigoplus_{l=1}^3 \mathbb{Q} \langle e_{(T_{a_i}^2)_l}^1 \times e_{(\mathcal{M}_{x_a})_i}^1 \rangle \bigoplus_{j=1}^b \bigoplus_{l=1}^2 \mathbb{Q} \langle e_{(T_{b_j}^2)_l}^1 \times e_{(\mathcal{M}_{x_b})_j}^1 \rangle \\
\mathcal{C}_1(\mathcal{L}_x) &= \bigoplus_{i=1}^a \mathbb{Q} \langle e_{(T_{a_i}^2)}^0 \times e_{(\mathcal{M}_{x_a})_i}^1 \rangle \bigoplus_{j=1}^b \mathbb{Q} \langle e_{(T_{b_j}^2)}^0 \times e_{(\mathcal{M}_{x_b})_j}^1 \rangle \\
\mathcal{C}_0(\mathcal{L}_x) &= \bigoplus_{l=0}^{f_{x_1}} \mathbb{Q} \langle e_{\mathcal{M}_{x_l}}^0 \rangle.
\end{aligned}$$

Remark 31. In the above chain complex, $e_{(\mathcal{M}_{x_a})_i}^1$ and $e_{(\mathcal{M}_{x_b})_j}^1$ are those 1-cells of \mathcal{M}_x , whose associated T^2 in \mathcal{L}_x has three and two 1-cells, respectively. Similarly, we have labeled each T^2 with either a_i or b_j according to the generator of the collapsed 1-dimensional cone in Σ .

Consequently, we obtain the following boundary operators for the above CW structure on \mathcal{L}_x .

$$\partial_5 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$(12) \quad \partial_4 = \begin{matrix} e_{T_1^3}^1 \times e_{\mathcal{M}_x}^2 \\ e_{T_2^3}^1 \times e_{\mathcal{M}_x}^2 \\ e_{T_3^3}^1 \times e_{\mathcal{M}_x}^2 \\ e_{(T_{a_i}^2)_1}^2 \times e_{(\mathcal{M}_{x_a})_i}^1 \\ e_{(T_{a_i}^2)_2}^2 \times e_{(\mathcal{M}_{x_a})_i}^1 \\ e_{(T_{b_j}^2)}^2 \times e_{(\mathcal{M}_{x_b})_j}^1 \end{matrix} \begin{matrix} a \\ a \\ a \\ a \\ a \\ b \end{matrix} \left[\begin{array}{ccc} e_{T_1^3}^2 \times e_{\mathcal{M}_x}^2 & e_{T_2^3}^2 \times e_{\mathcal{M}_x}^2 & e_{T_3^3}^2 \times e_{\mathcal{M}_x}^2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ n_{a_i} & m_{a_i} & l_{a_i} \\ n_{a_i} & m_{a_i} & l_{a_i} \\ n_{b_j} & m_{b_j} & 0 \end{array} \right]$$

$$(13) \quad \partial_3 = \begin{array}{c} \left. \begin{array}{l} e^0_{T^3} \times e^2_{\mathcal{M}_x} \\ e^1_{(T^2_{a_i})_1} \times e^1_{(\mathcal{M}_{x_a})_i} \\ e^1_{(T^2_{a_i})_2} \times e^1_{(\mathcal{M}_{x_a})_i} \\ e^1_{(T^2_{a_i})_3} \times e^1_{(\mathcal{M}_{x_a})_i} \\ e^1_{(T^2_{b_j})_1} \times e^1_{(\mathcal{M}_{x_b})_j} \\ e^1_{(T^2_{b_j})_2} \times e^1_{(\mathcal{M}_{x_b})_j} \end{array} \right\} \\ e^1_{T^3_1} \times e^2_{\mathcal{M}_x} \quad e^1_{T^3_2} \times e^2_{\mathcal{M}_x} \quad e^1_{T^3_3} \times e^2_{\mathcal{M}_x} \quad e^2_{(T^2_{a_i})_1} \times e^1_{(\mathcal{M}_{x_a})_i} \quad e^2_{(T^2_{a_i})_2} \times e^1_{(\mathcal{M}_{x_a})_i} \quad e^2_{(T^2_{b_j})} \times e^1_{(\mathcal{M}_{x_b})_j} \\ \left[\begin{array}{cccccc} 0 & 0 & 0 & \underbrace{0}_a & \underbrace{0}_a & \underbrace{0}_b \\ m_{a_i} l_{a_i} & 0 & 0 & 1 & -1 & 0 \\ 0 & n_{a_i} l_{a_i} & 0 & 1 & -1 & 0 \\ 0 & 0 & n_{a_i} m_{a_i} & 1 & -1 & 0 \\ -m_{b_j} & 0 & 0 & 0 & 0 & 0 \\ 0 & m_{b_j} & 0 & 0 & 0 & 0 \end{array} \right] \end{array}$$

In order to avoid any ambiguity on ∂_3 , we describe the explicit form of ∂_3 in more details. Let $l \in \{1, 2, 3\}$. The rows associated to $e^1_{(T_{a_i}^2)_l} \times e^1_{(\mathcal{M}_{x_a})_i}$ have only non-zero entries on the columns labeled by $e^1_{T_l^3} \times e^2_{\mathcal{M}_x}$, $e^2_{(T_{a_i}^2)_1} \times e^1_{(\mathcal{M}_{x_a})_i}$ and $e^2_{(T_{a_i}^2)_2} \times e^1_{(\mathcal{M}_{x_a})_i}$. Similarly, the columns labeled by $e^1_{T_l^3} \times e^2_{\mathcal{M}_x}$ have only non-zero entries on the rows labeled by $e^1_{(T_{a_i}^2)_l} \times e^1_{(\mathcal{M}_{x_a})_i}$ and possibly on $e^1_{(T_{b_j}^2)_1} \times e^1_{(\mathcal{M}_{x_b})_j}$ or $e^1_{(T_{b_j}^2)_2} \times e^1_{(\mathcal{M}_{x_b})_j}$. Lastly, the columns associated to $e^2_{(T_{a_i}^2)_1} \times e^1_{(\mathcal{M}_{x_a})_i}$ and $e^2_{(T_{a_i}^2)_2} \times e^1_{(\mathcal{M}_{x_a})_i}$ have only non-zero entries on the rows labeled by

$$e^1_{(T_{a_i}^2)_1} \times e^1_{(\mathcal{M}_{x_a})_i}, \quad e^1_{(T_{a_i}^2)_2} \times e^1_{(\mathcal{M}_{x_a})_i}, \quad \text{and} \quad e^1_{(T_{a_i}^2)_3} \times e^1_{(\mathcal{M}_{x_a})_i}.$$

For instance, let us start with $i = 1$. Consider the following part of ∂_3 :

$$\begin{pmatrix} m_{a_1} l_{a_1} & 0 & 0 & 1 & -1 & 0 \\ 0 & n_{a_1} l_{a_1} & 0 & 1 & -1 & 0 \\ 0 & 0 & n_{a_1} m_{a_1} & 1 & -1 & 0 \end{pmatrix}.$$

Adding $e^2_{(T_{a_i}^2)_1} \times e^1_{(\mathcal{M}_{x_a})_i}$ and $e^2_{(T_{a_i}^2)_2} \times e^1_{(\mathcal{M}_{x_a})_i}$ to columns, and $e^1_{(T_{a_i}^2)_1} \times e^1_{(\mathcal{M}_{x_a})_i}$, $e^1_{(T_{a_i}^2)_2} \times e^1_{(\mathcal{M}_{x_a})_i}$ and $e^1_{(T_{a_i}^2)_3} \times e^1_{(\mathcal{M}_{x_a})_i}$ to rows gives us

$$\begin{pmatrix} m_{a_1} l_{a_1} & 0 & 0 & 1 & -1 & 0 & 0 & 0 \\ 0 & n_{a_1} l_{a_1} & 0 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & n_{a_1} m_{a_1} & 1 & -1 & 0 & 0 & 0 \\ m_{a_2} l_{a_2} & 0 & 0 & 0 & 0 & 1 & -1 & 0 \\ 0 & n_{a_2} l_{a_2} & 0 & 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & n_{a_2} m_{a_2} & 0 & 0 & 1 & -1 & 0 \end{pmatrix}.$$

Adding more columns and rows goes along the same lines. Computing the \ker and Im of the above boundary operators is straightforward. We find

$$\begin{aligned} \text{rk}(\ker(\partial_5)) &= 1, \quad \text{rk}(\text{Im}(\partial_5)) = 0, \\ \text{rk}(\ker(\partial_4)) &= 0, \quad \text{rk}(\text{Im}(\partial_4)) = 3, \\ \text{rk}(\ker(\partial_3)) &= a + b = f_{x_1}, \end{aligned}$$

which yield

$$\begin{aligned} \text{rk}(H_5(\mathcal{L}_x; \mathbb{Q})) &= 1 \\ \text{rk}(H_4(\mathcal{L}_x; \mathbb{Q})) &= 0 \\ \text{rk}(H_3(\mathcal{L}_x; \mathbb{Q})) &= f_{x_1} - 3. \end{aligned}$$

The link \mathcal{L}_x is equipped with a stratification $\mathcal{L}_x = (\mathcal{L}_x)_5 \supset (\mathcal{L}_x)_3 \supset (\mathcal{L}_x)_1$ and a link of $(\mathcal{L}_x)_1$ in \mathcal{L}_x is a link of $(X_\Sigma)_2$ in X_Σ . As those links are rational homology spheres 11, \mathcal{L}_x is a rational homology manifold. In particular, the homology groups below the middle degree of \mathcal{L}_x can be obtained using rational Poincaré duality.

Remark 32. *There is yet another way for computing the homology groups below the middle degree. We can use our previous method of orthogonal decomposition and compute ∂_2 . The boundary operator ∂_2 will have the following form.*

$$\partial_2 = \begin{bmatrix} e_{T_1}^0 \times e_{\mathcal{M}_x}^2 & e_{(T_{a_i}^2)_1}^1 \times e_{(\mathcal{M}_{x_a})_i}^1 & e_{(T_{a_i}^2)_2}^1 \times e_{(\mathcal{M}_{x_a})_i}^1 & e_{(T_{a_i}^2)_3}^1 \times e_{(\mathcal{M}_{x_a})_i}^1 & e_{(T_{b_j}^2)_1}^1 \times e_{(\mathcal{M}_{x_b})_j}^1 & e_{(T_{b_j}^2)_2}^1 \times e_{(\mathcal{M}_{x_b})_j}^1 \\ \underbrace{e_{T_{a_i}^0}^0 \times e_{(\mathcal{M}_{x_a})_i}^1}_a & \underbrace{e_{T_{b_j}^0}^0 \times e_{(\mathcal{M}_{x_b})_j}^1}_b & \underbrace{e_{T_{a_i}^0}^0 \times e_{(\mathcal{M}_{x_a})_i}^1}_a & \underbrace{e_{T_{a_i}^0}^0 \times e_{(\mathcal{M}_{x_a})_i}^1}_a & \underbrace{e_{T_{b_j}^0}^0 \times e_{(\mathcal{M}_{x_b})_j}^1}_b & \underbrace{e_{T_{b_j}^0}^0 \times e_{(\mathcal{M}_{x_b})_j}^1}_b \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \underbrace{e_{S_i^1}^1 \times e_{(\mathcal{M}_x)_i}^{a+b}} & & & & \end{bmatrix}$$

It can be shown that rows associated to the cells of the form $e_{\mathcal{S}_i^1}^1 \times e_{(\mathcal{M}_x)_i}^0$ are linearly independent. This implies

$$\begin{aligned} \text{rk}(\ker(\partial_2)) &= (3a + 2b) - (a + b) = 2a + b \\ \text{rk}(\text{Im}(\partial_2)) &= a + b + 1. \end{aligned}$$

Keep in mind that the upper left part of ∂_2 is merely determined by the orientation of $|\mathcal{M}_x|$. Similarly, for ∂_1 we have

$$\partial_1 = e_{\mathcal{S}_i^1}^0 \times e_{(\mathcal{M}_x)_i}^0 \times e_{a+b} \left\{ \begin{bmatrix} e_{T_{a_i}^2}^0 \times e_{(\mathcal{M}_{x_a})_i}^1 & e_{T_{b_j}^2}^0 \times e_{(\mathcal{M}_{x_b})_j}^1 & e_{\mathcal{S}_i^1}^1 \times e_{(\mathcal{M}_x)_i}^0 \\ \underbrace{\quad}_a & \underbrace{\quad}_b & \underbrace{\quad}_{a+b} \\ z_i & y_j & 0 \end{bmatrix} \right\},$$

where again the y_j and the z_i are solely determined by the orientation of $|\mathcal{M}_x|$. Using $|\mathcal{M}_x| \cong \mathcal{D}^2$ yields

$$\begin{aligned} \text{rk}(\ker(\partial_1)) &= a + b + 1, \\ \text{rk}(\text{Im}(\partial_1)) &= f_{x_1} - 1, \\ \text{rk}(\ker(\partial_0)) &= f_{x_1}. \end{aligned}$$

In any case, we have shown the following proposition.

Proposition 33. *Let Σ be a complete 3-dimensional fan, X_Σ the associated toric variety and \mathcal{L}_x be the link of $x \in (X_\Sigma)_0$. Let f_{x_1} be the number of 1-dimensional faces of the dual polyhedron \mathcal{P} to Σ with x as a proper face. Then the Betti numbers of \mathcal{L}_x are given by*

$$\begin{aligned} \text{rk}(H_5(\mathcal{L}_x; \mathbb{Q})) &= 1, \\ \text{rk}(H_4(\mathcal{L}_x; \mathbb{Q})) &= 0, \\ \text{rk}(H_3(\mathcal{L}_x; \mathbb{Q})) &= f_{x_1} - 3, \\ \text{rk}(H_2(\mathcal{L}_x; \mathbb{Q})) &= f_{x_1} - 3, \\ \text{rk}(H_1(\mathcal{L}_x; \mathbb{Q})) &= 0, \\ \text{rk}(H_0(\mathcal{L}_x; \mathbb{Q})) &= 1. \end{aligned}$$

Corollary 34. *A point $x \in (X_\Sigma)_0$ is rationally singular if x in \mathcal{P} has more than three 1-dimensional neighboring faces.*

Proposition 35. *Let X_Σ be a 4-dimensional toric variety associated to a complete fan Σ with the stratification $X_\Sigma = X_4 \supset X_0$. Let $\nu \in X_0$ be a point whose link is an integral homology sphere. Then the link of ν , \mathcal{L}_ν , is in fact homeomorphic to \mathcal{S}^3 .*

Proof. Let \mathcal{P} be the dual polytope to Σ . Let $\sigma \in \Sigma$ be the cone dual to ν . The assumption on the link of ν implies that $H_1(\mathcal{L}_\nu; \mathbb{Z}) = 0$. Hence, the determinant of the two generators of σ vanishes, and they form a basis for \mathbb{Z}^2 . It follows that we can consider the images of the generators of σ under the projection map $\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2/\mathbb{Z}^2$ to be 1-cells of the minimal CW-structure of T^2 , which is attached to the interior of $\mathcal{I} \cong |\mathcal{M}_\nu|$. Let \mathcal{S}_1^1 and \mathcal{S}_2^1 be the image of the first and second generators of σ , respectively. Since the generators form a basis, we can set $T^2 \cong \mathcal{S}_1^1 \times \mathcal{S}_2^1$ to be the torus attached to $\text{int}(\mathcal{I})$. Let $p_{\mathcal{L}_\nu} : \mathcal{L}_\nu \rightarrow \mathcal{M}_\nu$ be the projection defined earlier. From the construction of the link, it follows that we have either $p_{\mathcal{L}_\nu}^{-1}(\{0\}) = \mathcal{S}_1^1$ and $p_{\mathcal{L}_\nu}^{-1}(\{1\}) = \mathcal{S}_2^1$ or $p_{\mathcal{L}_\nu}^{-1}(\{0\}) = \mathcal{S}_2^1$ and $p_{\mathcal{L}_\nu}^{-1}(\{1\}) = \mathcal{S}_1^1$. Hence, the link \mathcal{L}_ν is homeomorphic to the join $\mathcal{S}_1^1 * \mathcal{S}_2^1 \cong \mathcal{S}^3$, which proves our claim. \square

Proposition 36. *Let X_Σ be a 6-dimensional toric variety associated with a complete fan Σ with the stratification $X_\Sigma = X_6 \supset X_2 \supset X_0$. If $\nu \in X_2 - X_0$ is a point whose link is an integral homology sphere, then the link is in fact homeomorphic to S^3 . If $\nu \in X_0$ is a point whose link is an integral homology sphere, then the link of ν , \mathcal{L}_ν , is in fact homeomorphic to S^5 .*

Proof. If $\nu \in X_2 - X_0$, the statement follows from the proof of Proposition 35. Now, let $\nu \in X_0$ be a point whose link is an integral homology sphere. From the assumption, it follows that the base of the link, $|\mathcal{M}_\nu|$, is a 2-simplex. Let $\begin{pmatrix} n_1 \\ m_1 \\ l_1 \end{pmatrix}$, $\begin{pmatrix} n_2 \\ m_2 \\ l_2 \end{pmatrix}$, and $\begin{pmatrix} n_3 \\ m_3 \\ l_3 \end{pmatrix}$ be the generators of the dual cone to ν . For the moment, we assume that all entries are non-zero. Since $H_3(\mathcal{L}_\nu, \mathbb{Z}) = 0$, the image of the boundary operator

$$\partial_4 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ n_1 & m_1 & l_1 \\ n_1 & m_1 & l_1 \\ n_2 & m_2 & l_2 \\ n_2 & m_2 & l_2 \\ n_3 & m_3 & l_3 \\ n_3 & m_3 & l_3 \end{pmatrix}$$

is isomorphic to \mathbb{Z}^3 . It is the case if and only if

$$\det \begin{pmatrix} n_1 & m_1 & l_1 \\ n_2 & m_2 & l_2 \\ n_3 & m_3 & l_3 \end{pmatrix} = 1.$$

This means that the generators of the dual cone form a basis of \mathbb{Z}^3 . We attach the torus $T^3 \cong S^1 \times S^2 \times S^3$ to the interior of $|\mathcal{M}_\nu|$, where S^i is the image of $\begin{pmatrix} n_i \\ m_i \\ l_i \end{pmatrix}$ under the map $\mathbb{R}^3 \rightarrow \mathbb{R}^3/\mathbb{Z}^3$ for $i = 1, 2, 3$. On each 1-dimensional face of $|\mathcal{M}_\nu|$, we collapse one of the S^i in T^3 to a point. To each vertex of $|\mathcal{M}_\nu|$, we attach one of the S^i . From the construction, it follows that $\mathcal{L}_\nu \cong S^1 * S^3 \cong S^5$. We illustrate the above considerations in Figure 10.




FIGURE 10. The link of a smooth point $\nu \in X_0$ is homeomorphic to S^5

Now, let one of the generators have at least one zero entry. The boundary operator ∂_5 has five rows with at least one non-zero entry. In this case, the rest of the proof goes along the same lines as above. \square

5. \mathbb{Q} -HOMOLOGY PSEUDOMANIFOLDS, LEFSCHETZ DUALITY AND INTERSECTION HOMOLOGY

We have seen in previous sections that the link of a connected component of a 4-co-dimensional stratum in a toric variety is a rational homology 3-sphere. For instance, consider a 4-dimensional toric variety. As described earlier, the link of each isolated singularity is a rational homology 3-sphere. Such a toric variety satisfies Poincaré duality rationally. This suggests to consider strata with rational homology spherical links as homologically regular. We will thus introduce the notion of \mathbb{Q} -homology pseudomanifolds (or \mathbb{Q} -pseudomanifold for short), which is related to the concept of homology stratification considered by Rourke and Sanderson in [28].

Definition 37 (\mathbb{Q} -pseudomanifold). *We call a topological space X with filtration*

$$(14) \quad X \supset X_i \supset X_{i-1} \supset \cdots \supset X_0 \supset X_{-1} = \emptyset$$

a \mathbb{Q} -pseudomanifold if (14) can be augmented to a filtration of the form

$$X \supset X_{i+k} \supset \cdots \supset X_{i+1} \supset X_i \supset \cdots \supset X_0 \supset X_{-1} = \emptyset,$$

such that X is a pseudomanifold with respect to it and the link of each connected component of $X_{i+(j+1)} - X_{i+j}$ for $j = 0, \dots, k-1$ in X is a rational homology sphere. We call the filtration (14) a \mathbb{Q} -homology stratification (or \mathbb{Q} -stratification for short) of X .

Example 38. *A 6-dimensional toric variety X with $X = X_6 \supset X_0$ is a \mathbb{Q} -pseudomanifold, but generally not a stratified pseudomanifold with the respect to this stratification, since $X_6 - X_0$ is usually not a manifold, only a \mathbb{Q} -homology manifold.*

Definition 12 covers pseudomanifolds without boundary. We need to extend that definition to incorporate boundaries. The following definition is based on the corresponding PL-theoretic definition as formulated by Goresky and MacPherson in [22, Section 5.2, Page 155].

Definition 39 (Pseudomanifold with boundaries). *An n -dimensional stratified pseudomanifold with boundary is a pair (X, \mathcal{B}) of topological spaces such that \mathcal{B} is closed in X , together with a filtration $\{X_i\}$ on X such that:*

- (1) *The space $X - \mathcal{B}$ with the induced filtration $(X - \mathcal{B})_i = (X - \mathcal{B}) \cap X_i$ is an n -dimensional stratified pseudomanifold.*
- (2) *The space \mathcal{B} with the induced filtration $\mathcal{B}_{i-1} = \mathcal{B} \cap X_i$ is an $(n-1)$ -dimensional stratified pseudomanifold.*
- (3) *The topological space \mathcal{B} has an open filtered collar neighborhood in X , i.e. there exists an open neighborhood \mathcal{N} of \mathcal{B} and a filtered homeomorphism $\mathcal{N} \rightarrow [0, 1) \times \mathcal{B}$ (where $[0, 1)$ is given the trivial filtration) that takes \mathcal{B} to $\{0\} \times \mathcal{B}$ by the identity map.*

*The topological space \mathcal{B} is called the **boundary** of X and is also denoted by ∂X .*

The question arises to what extent the boundary \mathcal{B} is intrinsic to the topology of the space. The following proposition from Friedman-McClure [21] ensures that when there are no one-co-dimensional strata in X , the boundary \mathcal{B} depends only on the underlying space X and not the choice of a specific filtration (among those without one co-dimensional stratum).

Proposition 40. *Let $(X, \partial X)$ and $(X', \partial X')$ be equidimensional stratified pseudomanifolds with boundaries and no one-co-dimensional strata, and let $h : X \rightarrow X'$ be a homeomorphism (which is not required to be filtration preserving). Then h takes ∂X onto $\partial X'$.*

A **closed** pseudomanifold is a compact pseudomanifold with empty boundary.

In a 6-dimensional toric variety X , let \mathcal{C}_x be conical neighborhoods of $x \in X_0$, such that $\mathcal{C}_x \cap \mathcal{C}_{x'} = \emptyset$ for all $x, x' \in X_0$, $x \neq x'$. Removing somewhat smaller open cones in all \mathcal{C}_x from X gives us a pseudomanifold with boundary $(\mathcal{M}, \partial \mathcal{M})$, where now all links are rational homology spheres.

Definition 41. We call this process of removing a disjoint union of cone-like neighborhoods of X_0 **cutting out** the 0-dimensional stratum of X .

We shall prove sheaf-theoretically that compact oriented pseudomanifolds with boundary whose interior is a rational homology manifold satisfy Lefschetz duality for ordinary rational cohomology. We begin with some basic facts on direct and inverse limits. Let (I, \leq) be an (upwards) directed set, indexing direct systems $\{G_i\}_{i \in I}$, $\{H_i\}_{i \in I}$ of abelian groups. Let $\{f_i\} : \{G_i\} \rightarrow \{H_i\}$, $f_i : G_i \rightarrow H_i$, be a map of direct systems. If every f_i is an isomorphism, then

$$\lim_{\rightarrow I} f_i : \lim_{\rightarrow I} G_i \longrightarrow \lim_{\rightarrow I} H_i$$

is an isomorphism. (This is a consequence of the direct limit being a functor.) If $J \subset I$ is a cofinal subset, then the inclusion induces an isomorphism

$$\lim_{\rightarrow J} G_j \xrightarrow{\cong} \lim_{\rightarrow I} G_i$$

of direct limits. The diagram

$$\begin{array}{ccc} \lim_{\rightarrow J} G_j & \xrightarrow{\cong} & \lim_{\rightarrow I} G_i \\ \lim_{\rightarrow J} f_j \downarrow & & \downarrow \lim_{\rightarrow I} f_i \\ \lim_{\rightarrow J} H_j & \xrightarrow{\cong} & \lim_{\rightarrow I} H_i \end{array}$$

commutes. This has the following consequence. If every f_j , $j \in J$, is an isomorphism, then their direct limit is an isomorphism, and, by commutativity of the diagram, the direct limit of all f_i , $i \in I$, is an isomorphism as well. We make use of this principle at various points in the ensuing arguments.

Similar remarks apply to inverse limits. Let (I, \leq) be an (upwards) directed set, indexing inverse systems $\{G_i\}_{i \in I}$, $\{H_i\}_{i \in I}$ of abelian groups. Let $\{f_i\} : \{G_i\} \rightarrow \{H_i\}$, $f_i : G_i \rightarrow H_i$, be a map of inverse systems. If every f_i is an isomorphism, then

$$\lim_{\leftarrow I} f_i : \lim_{\leftarrow I} G_i \longrightarrow \lim_{\leftarrow I} H_i$$

is an isomorphism. (This is a consequence of the inverse limit being a functor.) If $J \subset I$ is a cofinal subset, then projection to components indexed by J induces an isomorphism

$$\lim_{\leftarrow I} G_i \xrightarrow{\cong} \lim_{\leftarrow J} G_j$$

of inverse limits. The diagram

$$\begin{array}{ccc} \lim_{\leftarrow I} G_i & \xrightarrow{\cong} & \lim_{\leftarrow J} G_j \\ \lim_{\leftarrow I} f_i \downarrow & & \downarrow \lim_{\leftarrow J} f_j \\ \lim_{\leftarrow I} H_i & \xrightarrow{\cong} & \lim_{\leftarrow J} H_j \end{array}$$

commutes. Again, we deduce that if every f_j , $j \in J$, is an isomorphism, then their inverse limit is an isomorphism, and, by commutativity of the diagram, the inverse limit of all f_i , $i \in I$, is an isomorphism as well.

The *cohomological dimension* of a topological space X over a commutative Noetherian ring R is the smallest $n \in \mathbb{N} \cup \{\infty\}$ such that $H_c^i(U; A) = 0$ for all open $U \subset X$, all sheaves A of R -modules on X , and all $i > n$. Topologically stratified pseudomanifolds of dimension n have cohomological dimension n (Borel [11, p. 60]). Recall that a topological space is called *first-countable* if each point has a countable neighborhood basis. Pseudomanifolds with boundary are first-countable, as can be seen by shrinking distinguished neighborhoods appropriately.

For the purposes of the present paper, we shall adopt the following concept of rational homology manifold with boundary.

Definition 42. A *rational homology n -manifold with boundary* is an n -dimensional topologically stratified pseudomanifold X with boundary ∂X such that $X - \partial X$ is a rational homology n -manifold. We will also refer to the pair $(X, \partial X)$ as a *\mathbb{Q} -manifold with boundary*.

In particular, rational homology manifolds $(X, \partial X)$ with boundary in the sense of this definition are paracompact Hausdorff spaces which are locally compact and (strongly) locally contractible. (A space is *strongly locally contractible* if every point possesses a neighborhood basis consisting of contractible sets.) Since they are topologically stratified pseudomanifolds, rational homology manifolds with boundary have finite cohomological dimension over \mathbb{Q} , and their cohomological dimension agrees with their pseudomanifold dimension. Note that in the above definition, we do not explicitly require ∂X to be a rational homology $(n - 1)$ -manifold. This turns out to be true, as we shall see in Proposition 50.

The cohomology sheaf in degree $k \in \mathbb{Z}$ of a differential graded sheaf S on a space X will be denoted by $\mathbf{H}^k(S)$, the hypercohomology by $\mathcal{H}^k(X; S)$. If S is concentrated in degree 0, then $\mathcal{H}^k(X; S)$ is the sheaf cohomology group $H^k(X; S)$. The constant sheaf with stalk \mathbb{Q} on a space X will be written as \mathbb{Q}_X and will frequently be considered as a differential graded sheaf concentrated in degree 0.

Lemma 43. Let $(X, \partial X)$ be a pseudomanifold with boundary and let i be the open inclusion $i : X - \partial X \hookrightarrow X$. Then the canonical adjunction map

$$\mathbb{Q}_X \longrightarrow Ri_*i^*\mathbb{Q}_X$$

is a quasi-isomorphism.

Proof. We recall that the property of being a quasi-isomorphism can be checked point by point on cohomology stalks. Let S be any differential graded sheaf on X . The adjunction $S \rightarrow Ri_*i^*S$ restricts to the identity $i^*S \rightarrow i^*S$ on $X - \partial X$ and is therefore a quasi-isomorphism at every point $x \in X - \partial X$. It remains to check that it is a quasi-isomorphism at points x in ∂X .

Let U be an open neighborhood of $x \in \partial X$ in X . By definition of a pseudomanifold with boundary, ∂X is collared in X . Thus there exists an open neighborhood N of ∂X and a (filtered) homeomorphism $\phi : N \rightarrow [0, 1) \times \partial X$ which restricts to the identity $\partial X \rightarrow \{0\} \times \partial X$. Note that ϕ restricts to a homeomorphism $\phi : N - \partial X \rightarrow (0, 1) \times \partial X$. The intersection $U \cap N$ is an open neighborhood of x in X . Thus $\phi(U \cap N)$ is an open neighborhood of $\phi(x) = (0, x)$ in $[0, 1) \times \partial X$. By definition of the product topology, there exists an open neighborhood $V \subset \partial X$ of x in ∂X and $\epsilon > 0$ such that $[0, \epsilon) \times V \subset \phi(U \cap N)$. Set

$$U' := \phi^{-1}([0, \epsilon) \times V).$$

Then $U' \subset X$ is an open neighborhood of $\phi^{-1}(0, x) = x$ in X and $U' \subset U$, for

$$U' = \phi^{-1}([0, \epsilon) \times V) \subset \phi^{-1}(\phi(U \cap N)) = U \cap N \subset U.$$

We have thus shown that every point $x \in \partial X$ has a neighborhood basis consisting of open sets of the form U' . We note that $U' - \partial X \subset U - \partial X$.

The sheafification \mathbf{A} of a presheaf $U \mapsto A(U)$ has stalks

$$\mathbf{A}_x = \varinjlim A(U),$$

where the direct limit ranges over all open neighborhoods U of x . Since, by definition, $\mathbf{H}^k(S)$ is the sheafification of the presheaf $U \mapsto H^k\Gamma(U; S)$ (Iversen [24, p. 89]), the cohomology sheaf has stalks

$$\mathbf{H}^k(S)_x = \varinjlim H^k\Gamma(U; S).$$

We note that $\mathbf{H}^k(S)_x = H^k(S_x)$, as restriction to stalks is an exact functor. Thus for the constant sheaf $S = \mathbb{Q}_X$ on X ,

$$\mathbf{H}^0(\mathbb{Q}_X)_x = H^0(\mathbb{Q}_{X,x}) = \mathbb{Q}$$

and

$$\mathbf{H}^k(\mathbb{Q}_X)_x = H^k(\mathbb{Q}_{X,x}) = 0, \quad k \neq 0.$$

An injective resolution $S \rightarrow I$ of S is a quasi-isomorphism and thus induces an isomorphism $\mathbf{H}^k(S) \cong \mathbf{H}^k(I)$. Therefore, the cohomology stalks may also be computed as

$$\mathbf{H}^k(S)_x = \varinjlim H^k\Gamma(U; I).$$

Now the groups $H^k\Gamma(U; I)$ are precisely the hypercohomology groups $\mathcal{H}^k(U; S)$. We obtain the formula

$$\mathbf{H}^k(S)_x = \varinjlim \mathcal{H}^k(U; S).$$

The restriction of an injective sheaf to an open subset is injective ([24, p. 109, Cor. 6.10]). Therefore, $i^*S \rightarrow i^*I$ is an injective resolution of i^*S . We deduce that $Ri_*i^*S = i_*i^*I$ and the canonical adjunction morphism $S \rightarrow Ri_*i^*S$ is given by the composition

$$S \longrightarrow I \longrightarrow i_*i^*I = Ri_*i^*S.$$

The cohomology stalks of Ri_*i^*S are given by

$$\begin{aligned} \mathbf{H}^k(Ri_*i^*S)_x &= \mathbf{H}^k(i_*i^*I)_x = \varinjlim H^k\Gamma(U; i_*i^*I) \\ &= \varinjlim H^k\Gamma(U - \partial X; i^*I) = \varinjlim \mathcal{H}^k(U - \partial X; S). \end{aligned}$$

At $x \in \partial X$, the map

$$\mathbf{H}^k(S)_x \longrightarrow \mathbf{H}^k(Ri_*i^*S)_x$$

is hence given by the map

$$\varinjlim \mathcal{H}^k(U; S) \longrightarrow \varinjlim \mathcal{H}^k(U - \partial X; S)$$

induced on direct limits by the restriction maps

$$(15) \quad \mathcal{H}^k(U; S) \longrightarrow \mathcal{H}^k(U - \partial X; S).$$

Since an arbitrary neighborhood U contains one of the form U' , neighborhoods of the latter type are cofinal in the directed set of all open neighborhoods. So the direct limit can be computed on neighborhoods of type U' . When $S = \mathbb{Q}_X$ is the constant sheaf, the collar homeomorphism ϕ identifies the restriction map

$$\mathcal{H}^k(U'; \mathbb{Q}_X) \longrightarrow \mathcal{H}^k(U' - \partial X; \mathbb{Q}_X)$$

with the restriction map

$$(16) \quad \mathcal{H}^k([0, \epsilon) \times V; \mathbb{Q}_X) \longrightarrow \mathcal{H}^k((0, \epsilon) \times V; \mathbb{Q}_X).$$

Consider the commutative diagram

$$\begin{array}{ccc} (0, \epsilon) \times V & \xrightarrow{\quad} & [0, \epsilon) \times V \\ & \searrow & \swarrow \\ & V & \end{array}$$

where the maps to V are the second factor projections. These projections have contractible fibers and are homotopy equivalences. It follows that the horizontal inclusion is a homotopy equivalence. On singular cohomology, which is homotopy invariant, we get an induced restriction isomorphism

$$(17) \quad H^k([0, \epsilon) \times V; \mathbb{Q}) \xrightarrow{\cong} H^k((0, \epsilon) \times V; \mathbb{Q}).$$

For semi-locally contractible topological spaces, there is a natural isomorphism between singular cohomology $H^*(-; \mathbb{Q})$ and sheaf cohomology $H^*(-; \mathbb{Q}_X)$ (Sella [30]). Locally contractible spaces are in particular semi-locally contractible. The space $[0, \epsilon) \times V$ is locally contractible, since V is an open subset of the locally contractible space ∂X . The natural isomorphism between singular and sheaf cohomology thus identifies the maps (17) and (16). Therefore, the latter is an isomorphism as well. The basic principle on direct limits recalled earlier implies that the direct limit of the maps (15) is an isomorphism. \square

We shall use the term *cohomologically constructible*, a property of differential graded sheaves, in the sense of Borel [11, p. 69]. We will not recall the definition here, but point out that this notion does not require a stratification. According to [11, p. 79, Cor. 3.11 (i)], the constant sheaf on a stratified pseudomanifold without boundary is cohomologically constructible. We need to extend this statement to pseudomanifolds with boundary:

Lemma 44. *Let $(X, \partial X)$ be a topologically stratified pseudomanifold with boundary. Then the constant sheaf \mathbb{Q}_X on X is cohomologically constructible.*

Proof. Since the interior $X - \partial X$ is a pseudomanifold without boundary, \mathbb{Q}_X is cohomologically constructible at every point of the interior by [11, p. 79, Cor. 3.11 (i)]. It remains to verify that \mathbb{Q}_X is cohomologically constructible at boundary points $x \in \partial X$. Using a collar as in the proof of Lemma 43, there exists a neighborhood basis $\{U_i\}_{i \in I}$ of x in X consisting of open sets U_i that are homeomorphic to $[0, \epsilon_i) \times V_i$, $V_i \subset \partial X$, V_i open and contractible, $\epsilon_i > 0$. If $i < j$ so that $U_i \supset U_j$, then the restriction map

$$\mathcal{H}^k(U_i; \mathbb{Q}_X) \longrightarrow \mathcal{H}^k(U_j; \mathbb{Q}_X)$$

is an isomorphism, since in the commutative diagram

$$\begin{array}{ccc} U_j & \xrightarrow{\subset} & U_i \\ & \searrow & \swarrow \\ & \text{pt} & \end{array}$$

the constant maps to the point are homotopy equivalences, and thus the horizontal inclusion is a homotopy equivalence. (Here, as earlier, we may identify singular and sheaf cohomology.) Thus, the direct system $\{\mathcal{H}^k(U_i; \mathbb{Q}_X)\}_{i \in I}$ is constant, hence also essentially constant in the sense of Borel's definition [11, p. 68]. Since $\{U_i\}_{i \in I}$ is cofinal in the directed set of all open neighborhoods of x , the direct system $\{\mathcal{H}^k(U; \mathbb{Q}_X)\}$ over all neighborhoods of x is also essentially constant ([11, p. 69]). We observe furthermore that the direct limit is finitely generated as it is one-dimensional for $k = 0$ and 0 otherwise. This proves condition CC2 of Borel.

We shall next treat the dual case of inverse systems given by compactly supported cohomology. Thus we consider the inverse system with groups $\mathcal{H}_c^k(U_i; \mathbb{Q}_X)$, $i \in I$, where $\{U_i\}$ is the same neighborhood basis of $x \in \partial X$ as before. For $i < j$, so that $U_i \supset U_j$, the transition maps are given by extension by zero

$$\mathcal{H}_c^k(U_j; \mathbb{Q}_X) \longrightarrow \mathcal{H}_c^k(U_i; \mathbb{Q}_X).$$

We claim that in fact all of these groups vanish. Indeed, we may use the Künneth formula

$$R\Gamma_c^\bullet(A \times B; \mathbb{Q}) = R\Gamma_c^\bullet(A; \mathbb{Q}) \otimes_{\mathbb{Q}} R\Gamma_c^\bullet(B; \mathbb{Q})$$

valid for finite dimensional locally compact spaces A, B , [24, p. 323]. The spaces $[0, \epsilon_i)$ and V_i satisfy these assumptions and therefore

$$\mathcal{H}_c^*(U_i; \mathbb{Q}_X) \cong \mathcal{H}_c^*([0, \epsilon_i) \times V_i; \mathbb{Q}_X) \cong \mathcal{H}_c^*([0, \epsilon_i); \mathbb{Q}) \otimes \mathcal{H}_c^*(V_i; \mathbb{Q}).$$

Now, the compactly supported cohomology with constant coefficients of a closed halfspace vanishes, [24, p. 189, 8.4]. So $\mathcal{H}_c^*([0, \epsilon_i); \mathbb{Q}) = 0$ and hence $\mathcal{H}_c^*(U_i; \mathbb{Q}_X) = 0$ as claimed. The inverse

system $\{\mathcal{H}_c^k(U_i; \mathbb{Q}_X)\}_{i \in I}$ is thus constant, hence also essentially constant. Since $\{U_i\}_{i \in I}$ is cofinal in the directed set of all open neighborhoods of x , the inverse system $\{\mathcal{H}_c^k(U; \mathbb{Q}_X)\}$ over all open neighborhoods of x is also essentially constant ([11, p. 69]). The inverse limit is zero, so in particular finitely generated. This proves condition CC1 of Borel.

We have verified Borel's conditions CC1 and CC2. These imply the remaining conditions CC3 and CC4, as pointed out by Borel. \square

Let $(X, \partial X)$ be a rational homology n -manifold with boundary. By definition, $X - \partial X$ is a rational homology n -manifold (without boundary). Thus the dualizing complex $\mathbb{D}_{X-\partial X}[-n]$ is naturally quasi-isomorphic to the orientation sheaf $\text{or}_{X-\partial X}$ of $X - \partial X$.

Definition 45. *The homology manifold $(X, \partial X)$ with boundary is called **orientable** if there exists an isomorphism $\text{or}_{X-\partial X} \cong \mathbb{Q}_{X-\partial X}$. A choice of such an isomorphism, if it exists, is called an **orientation**.*

Let $(X, \partial X)$ be an oriented rational homology n -manifold with boundary. The orientation induces a quasi-isomorphism

$$\mathbb{D}_{X-\partial X}[-n] \cong \text{or}_{X-\partial X} \cong \mathbb{Q}_{X-\partial X}.$$

Using Verdier's dualizing functor \mathcal{D} , we may express the dualizing complex as $\mathbb{D}_{X-\partial X} = \mathcal{D}\mathbb{Q}_{X-\partial X}$ ([23, p. 90]). The orientation therefore yields a self-duality isomorphism

$$(\mathcal{D}\mathbb{Q}_{X-\partial X})[-n] \cong \mathbb{Q}_{X-\partial X}.$$

Lemma 46. *(Borel [11, p. 137, Prop. 8.8 (3)].) Let Y be a first-countable locally compact space. Let $i : U \subset Y$ be an open inclusion and let B be a differential graded sheaf on U such that Ri_*B is cohomologically constructible. Then*

$$\mathcal{D}Ri_*B = i_!\mathcal{D}B.$$

We apply this lemma to $i : X - \partial X \hookrightarrow X$, observing that X is locally compact and first-countable. We take $B = \mathbb{Q}_{X-\partial X}$. By Lemma 43,

$$Ri_*B = Ri_*i^*\mathbb{Q}_X \cong \mathbb{Q}_X,$$

the constant sheaf on X , which is cohomologically constructible by Lemma 44. By Lemma 46,

$$\mathcal{D}Ri_*\mathbb{Q}_{X-\partial X} = i_!\mathcal{D}\mathbb{Q}_{X-\partial X}.$$

Using the above self-duality isomorphism,

$$\begin{aligned} i_!i^*\mathbb{Q}_X &= i_!\mathbb{Q}_{X-\partial X} \cong i_!\mathcal{D}\mathbb{Q}_{X-\partial X}[-n] \\ &= \mathcal{D}Ri_*\mathbb{Q}_{X-\partial X}[-n] = \mathcal{D}Ri_*i^*\mathbb{Q}_X[-n] \\ &= \mathcal{D}\mathbb{Q}_X[-n]. \end{aligned}$$

On hypercohomology, there is an induced isomorphism

$$\mathcal{H}^k(X; i_!i^*\mathbb{Q}_X) \cong \mathcal{H}^{k-n}(X; \mathcal{D}\mathbb{Q}_X).$$

The global effect of dualizing a differential graded sheaf is

$$\mathcal{H}^{k-n}(X; \mathcal{D}\mathbb{Q}_X) = \text{Hom}(\mathcal{H}^{n-k}(X; \mathbb{Q}_X), \mathbb{Q}).$$

(This holds on any locally compact space which has finite cohomological dimension over \mathbb{Q} .) For compact X , $\mathcal{H}_c^{n-k}(X; \mathbb{Q}_X) = \mathcal{H}^{n-k}(X; \mathbb{Q}_X)$, so we obtain an isomorphism

$$\mathcal{H}^{k-n}(X; \mathcal{D}\mathbb{Q}_X) = \text{Hom}(\mathcal{H}^{n-k}(X; \mathbb{Q}_X), \mathbb{Q}).$$

Composing, we receive a duality isomorphism

$$\mathcal{H}^k(X; i_!i^*\mathbb{Q}_X) \cong \text{Hom}(\mathcal{H}^{n-k}(X; \mathbb{Q}_X), \mathbb{Q})$$

for compact X . The left hand group is nothing but the relative group $H^k(X, \partial X; \mathbb{Q})$ (Iversen [24, p. 249]), while the universal coefficient theorem identifies the right hand group with the homology group $H_{n-k}(X; \mathbb{Q})$. The resulting isomorphism

$$H^k(X, \partial X; \mathbb{Q}) \cong H_{n-k}(X; \mathbb{Q})$$

is Lefschetz duality for an oriented compact rational homology n -manifold $(X, \partial X)$ with boundary. We have proved:

Theorem 47. *Let $(X, \partial X)$ be an oriented compact rational homology n -manifold with boundary. Then $(X, \partial X)$ has a Lefschetz duality isomorphism*

$$H^k(X, \partial X; \mathbb{Q}) \cong H_{n-k}(X; \mathbb{Q}).$$

Corollary 48. *Let X be an n -dimensional compact oriented \mathbb{Q} -pseudomanifold with \mathbb{Q} -stratification*

$$X = X_n \supset X_0.$$

Let $(\mathcal{M}, \partial\mathcal{M})$ be the pseudomanifold with boundary obtained by cutting out the 0-dimensional stratum in the \mathbb{Q} -stratification. Then $(\mathcal{M}, \partial\mathcal{M})$ satisfies Lefschetz duality with respect to ordinary rational homology.

Proof. The pair $(\mathcal{M}, \partial\mathcal{M})$ is a compact oriented rational homology manifold with boundary. The result follows from Theorem 47. \square

Corollary 49. *Let X be an n -dimensional closed oriented \mathbb{Q} -pseudomanifold with \mathbb{Q} -stratification $X = X_n$. (Such an X is a rational homology manifold.) Then X satisfies Poincaré duality rationally.*

Proposition 50. *If $(X, \partial X)$ is a rational homology n -manifold with boundary, then ∂X is a rational homology $(n-1)$ -manifold (without boundary).*

Proof. Sheaf-theoretically, this is a direct consequence of [3, Chapter 4] and Lemma 43. The subspace ∂X has a stratum preserving collar neighborhood in X . This implies that links in ∂X can be taken to be links in $X - \partial X$. \square

Example 51. *Let X be a 4-dimensional toric variety. Then X satisfies Poincaré duality rationally.*

Example 52. *Let X be a 6-dimensional toric variety. The pseudomanifold obtained by cutting out X_0 satisfies Lefschetz duality, rationally.*

Concerning intersection homology, we shall use the following standard notation.

Definition 53. (Goresky-MacPherson.) *A **perversity***

$$\bar{p} : \mathbb{Z}_{\geq 2} \longrightarrow \mathbb{Z}$$

*is a function such that $\bar{p}(2) = 0$ and $\bar{p}(k+1) - \bar{p}(k) \in \{1, 0\}$. The **complementary perversity** \bar{q} of \bar{p} is the one with $\bar{p}(k) + \bar{q}(k) = k - 2$.*

We denote the i -th intersection homology group of X with coefficients in A and the perversity \bar{p} with $I^{\bar{p}}H_i(X; A)$. Similarly, for the i -th intersection cohomology group, we write $I_{\bar{p}}H^i(X; A)$. For an introduction to the theory of intersection homology, the reader may consult [20] by Friedman, [25] by Kirwan and Woolf, or [4].

6. INTERSECTION SPACES AND ISOLATED SINGULARITIES

In this section, first, we briefly recall the theory of intersection spaces introduced by the first named author in [6]. For toric varieties, we only consider the duality of Betti numbers for the middle perversity. Throughout this section, we work with rational coefficients unless otherwise specified.

Definition 54. *Let n be a natural number. A CW complex \mathcal{K} is called **rationally n -segmented** if it contains a sub-complex $\mathcal{K}_{<n} \subset \mathcal{K}$ such that*

$$H_r(\mathcal{K}_{<n}) = 0 \text{ for } r \geq n \text{ and } i_* : H_r(\mathcal{K}_{<n}) \xrightarrow{\cong} H_r(\mathcal{K}) \text{ for } r < n,$$

where i is the inclusion of $\mathcal{K}_{<n}$ into \mathcal{K} .

Given any CW complex \mathcal{K} and natural number n , there exists a **homology n -truncation** (Moore approximation) $f : \mathcal{K}_{<n} \rightarrow \mathcal{K}$, i.e. a continuous map from a CW complex $\mathcal{K}_{<n}$ to \mathcal{K} such that $f_* : H_*(\mathcal{K}_{<n}; \mathbb{Z}) \rightarrow H_*(\mathcal{K}; \mathbb{Z})$ is an isomorphism for $* < n$ and $H_*(\mathcal{K}_{<n}; \mathbb{Z}) = 0$ for $* \geq n$, see [6] (for the simply connected case) and Wrazidlo [34] (in general). The map f cannot in general be taken to be a sub-complex inclusion. We shall prove in Proposition 67 that the 5-dimensional links in a 6-dimensional toric variety are rationally 3-segmented.

Let $X \supset X_0$ be an n -dimensional compact oriented topological pseudomanifold with isolated singularities. For $x_i \in X_0$, let \mathcal{L}_i be an associated link. Although two different links associated with the same isolated singularity need not be homeomorphic, we shall justify that the homology of links is well-defined. Let \mathcal{L}_i be as above and \mathcal{U} an open distinguished neighborhood of x_i . Then there exists a homeomorphism $\phi : \mathcal{U} \xrightarrow{\cong} \mathcal{C}(\mathcal{L})$ which sends x_i to the cone point, c . We define $\mathcal{C}_{1/2}(\mathcal{L}_i) \subset \mathcal{C}(\mathcal{L}_i)$ as the open cone over \mathcal{L}_i by considering the interval $[0, \frac{1}{2}] \subset [0, 1]$. Let $\mathcal{U}_{1/2} \subset \mathcal{U}$ be the preimage of $\mathcal{C}_{1/2}(\mathcal{L}_i)$ under the homeomorphism ϕ . By employing the excision axiom, we have $H_*(X, X - x_i) \cong H_*(X - (X - \mathcal{U}_{1/2}), (X - x_i) - (X - \mathcal{U}_{1/2})) = H_*(\mathcal{U}_{1/2}, \mathcal{U}_{1/2} - x_i)$. By using the homeomorphism ϕ we arrive at $H_*(X, X - x_i) \cong H_*(\mathcal{C}_{1/2}(\mathcal{L}_i), \mathcal{C}_{1/2}(\mathcal{L}_i) - \{c\})$. As $\mathcal{C}_{1/2}(\mathcal{L}_i) \simeq *$ and $\mathcal{C}_{1/2}(\mathcal{L}_i) - \{c\} \simeq \mathcal{L}_i$, the connecting homomorphism in the long exact sequence of the relative homology groups of the pair $(\mathcal{C}_{1/2}(\mathcal{L}_i), \mathcal{C}_{1/2}(\mathcal{L}_i) - \{c\})$ is an isomorphism. If \mathcal{L}'_i is another link of x_i , then the above procedure yields $H_*(\mathcal{L}_i) \cong H_*(\mathcal{L}'_i)$. Hence the homology of links is well-defined.

Assume that all links \mathcal{L}_i can be equipped with CW structures such that they are rationally k -segmented, where $k = n - 1 - \bar{p}(n)$, for the perversity \bar{p} . Let $(\mathcal{M}, \partial\mathcal{M})$ be the manifold with boundary obtained by cutting out all isolated singularities of X . Let $(\mathcal{L}_i)_{<k}$ be a sub-complex of \mathcal{L}_i that truncates the homology. Then, we have $\partial\mathcal{M} = \bigsqcup_i \mathcal{L}_i$. Let $\mathcal{L}_{<k} = \bigsqcup_i (\mathcal{L}_i)_{<k}$ and define a homotopy class g by the composition $g : \mathcal{L}_{<k} \hookrightarrow \partial\mathcal{M} \xrightarrow{\text{incl.}} \mathcal{M}$. For the purposes of the present paper we adopt the following definition:

Definition 55. *The perversity \bar{p} rational intersection space $I^{\bar{p}}X$ of X is defined to be*

$$I^{\bar{p}}X = \text{cone}(g) = \mathcal{M} \bigcup_g \mathcal{C}(\mathcal{L}_{<k}).$$

Due to the use of rational coefficients, $I^{\bar{p}}X$ as defined above need not be integrally homotopy equivalent to the construction of [6], but the rational homology groups are isomorphic.

Theorem 56. *Let X be an n -dimensional compact oriented topological pseudomanifold with only isolated singularities. Let \bar{p} and \bar{q} be complementary perversities. Then, we have the duality isomorphism*

$$d : \tilde{H}_r(I^{\bar{p}}X)^* \xrightarrow{\cong} \tilde{H}_{n-r}(I^{\bar{q}}X),$$

where

$$\tilde{H}_r(I^{\bar{p}}X)^* = \text{Hom}(\tilde{H}_r(I^{\bar{p}}X), \mathbb{Q}).$$

Proof. The detailed construction of the above duality isomorphism can be found in the proof of [6, Theorem 2.12] by the first named author. However, as mentioned, $I^{\bar{p}}X$, as defined above need not be homotopy equivalent to the construction of [6]. Hence, we need to adjust the previous proof slightly. In our setting we have the inclusion $\mathcal{L}_{<k} \hookrightarrow \mathcal{L} = \partial\mathcal{M}$. We consider the same braid as in the proof in loc. cit. One can then proceed along the same line as in [6]. \square

Remark 57. *Let $k = n - 1 - \bar{p}(n)$. Then, we have the following isomorphisms. For $r > k$,*

$$\begin{aligned} H_r(\mathcal{M}) &\xrightarrow{\cong} \tilde{H}_r(I^{\bar{p}}X) \\ \tilde{H}_{n-r}(I^{\bar{q}}X) &\xrightarrow{\cong} H_{n-r}(\mathcal{M}, \partial\mathcal{M}). \end{aligned}$$

For $r < k$,

$$\begin{aligned} \tilde{H}_r(I^{\bar{p}}X) &\xrightarrow{\cong} H_r(\mathcal{M}, \partial\mathcal{M}) \\ H_{n-r}(\mathcal{M}) &\xrightarrow{\cong} \tilde{H}_{n-r}(I^{\bar{q}}X). \end{aligned}$$

Corollary 58. *If $n = \dim(X)$ is even, then the difference between the Euler characteristic of $\tilde{H}_*(I^{\bar{p}}X)$ and $I^{\bar{p}}H_*(X)$ is given by*

$$\chi(\tilde{H}_*(I^{\bar{p}}X)) - \chi(I^{\bar{p}}H_*(X)) = -2\chi_{<n-1-\bar{p}(n)}(\mathcal{L}),$$

where \mathcal{L} is the disjoint union of the links of the isolated singularities of X . If $n = \dim(X)$ is odd, then

$$\chi(\tilde{H}_*(I^{\bar{n}}X)) - \chi(I^{\bar{n}}H_*(X)) = (-1)^{\frac{n-1}{2}} b_{\frac{n-1}{2}}(\mathcal{L}),$$

where $b_{\frac{n-1}{2}}(\mathcal{L})$ is the middle dimensional Betti number of \mathcal{L} and \bar{n} is the upper middle perversity. Regardless of the parity of n , the identity

$$\mathrm{rk}(\tilde{H}_k(I^{\bar{p}}X)) + \mathrm{rk}(I^{\bar{p}}H_k(X)) = \mathrm{rk}(H_k(\mathcal{M})) + \mathrm{rk}(H_k(\mathcal{M}, \mathcal{L}))$$

always holds in degree $k = n - 1 - \bar{p}(n)$, where \mathcal{M} is the exterior of the singular set of X .

Proof. A proof of the above corollary can be found in [6] by the first named author. \square

Lemma 59. *The rational homology of $I^{\bar{p}}X$ is independent of the choice of $(\mathcal{L}_i)_{<k}$.*

Proof. From Remark 57 it follows immediately that $H_r(I^{\bar{p}}X)$ does not depend of the choice of $(\mathcal{L}_i)_{<k}$ for $r < k$ and $r > k$. For the case $r = k$ the claim follows from the last identity in Corollary 58. \square

7. CONSTRUCTION OF INTERSECTION SPACES FOR \mathbb{Q} -PSEUDOMANIFOLDS

In this section, we generalize the theory of intersection spaces to \mathbb{Q} -pseudomanifolds with \mathbb{Q} -isolated singularities. To do this, we revisit the proof of Theorem 56 introduced by the first named author in [6]. The central idea is to indicate that the main ingredients of that proof remain intact, even for \mathbb{Q} -pseudomanifold with isolated singularities. Before proving our main theorem, we want to compute the intersection homology of \mathbb{Q} -pseudomanifolds with \mathbb{Q} -isolated singularities. But first of all, we need to generalize the previous definition of intersection spaces to the \mathbb{Q} -isolated setting.

Let X be an n -dimensional compact oriented \mathbb{Q} -pseudomanifold with \mathbb{Q} -isolated singularities, which means X has a \mathbb{Q} -stratification of the form

$$X = X_n \supset X_0.$$

For $x_i \in X_0$, let \mathcal{L}_i be an associated link. Assume that all links are rationally k -segmented, where $k = n - 1 - \bar{p}(n)$, for the perversity \bar{p} . Let \mathcal{M} be the compact \mathbb{Q} -manifold with boundary

obtained by cutting out all \mathbb{Q} -isolated singularities of X . Then, we have $\partial\mathcal{M} = \bigsqcup_i \mathcal{L}_i$. Let $\mathcal{L}_{<k} = \bigsqcup_i (\mathcal{L}_i)_{<k}$, and define a homotopy class $g : \mathcal{L}_{<k} \hookrightarrow \partial\mathcal{M} \xrightarrow{\text{incl.}} \mathcal{M}$.

Definition 60. *The perversity \bar{p} **generalized intersection space** $I^{\bar{p}}X$ of X is defined to be*

$$I^{\bar{p}}X = \text{cone}(g) = \mathcal{M} \bigcup_g \mathcal{C}(\mathcal{L}_{<k}).$$

We shall use the standard cone computation for intersection homology:

Proposition 61. *Suppose Y is a compact topological pseudomanifold of dimension $m - 1 \geq 0$. Then for a perversity \bar{p} ,*

$$I^{\bar{p}}H_r(\mathcal{C}(Y)) = \begin{cases} I^{\bar{p}}H_r(Y), & r < m - 1 - \bar{p}(m) \\ 0, & \text{otherwise.} \end{cases}$$

Proposition 62. *Let X be an n -dimensional \mathbb{Q} -pseudomanifold with \mathbb{Q} -isolated singularities, where $n \geq 1$. Then for perversity \bar{p} , we have*

$$I^{\bar{p}}H_r(X) \cong \begin{cases} H_r(\mathcal{M}, \mathcal{L}), & r > k \\ H_r(\mathcal{M}), & r < k, \end{cases}$$

where \mathcal{M} is again the \mathbb{Q} -manifold obtained by cutting out the \mathbb{Q} -isolated singularities and $\mathcal{L} = \bigsqcup_i \mathcal{L}_{x_i}$, where for each $x_i \in X_0$, \mathcal{L}_{x_i} is a link associated to x_i , and $k = n - 1 - \bar{p}(n)$.

Proof. We cut out cone-like neighborhoods of \mathbb{Q} -isolated singularities. The obtained topological space is a pseudomanifold with boundary, $(\mathcal{M}, \partial\mathcal{M})$. In fact $(\mathcal{M}, \partial\mathcal{M})$ is a \mathbb{Q} -manifold with boundary. Let $\mathcal{C}(\mathcal{L}_{x_i})$ be the removed open cone-like neighborhood of x_i . Let \mathcal{V} be an open neighborhood of \mathcal{M} in X that deformation retracts in a stratum preserving way to \mathcal{M} and set $\mathcal{U} = \bigsqcup_i \mathcal{C}(\mathcal{L}_{x_i})$. Thus, we have $\cap := \mathcal{U} \cap \mathcal{V} \cong \bigsqcup_i \mathcal{L}_{x_i} \times (\frac{1}{2}, 1)$. The Mayer-Vietoris sequence for intersection homology is a long exact sequence

$$\dots \longrightarrow I^{\bar{p}}H_r(\cap) \longrightarrow I^{\bar{p}}H_r(\mathcal{U}) \oplus I^{\bar{p}}H_r(\mathcal{V}) \longrightarrow I^{\bar{p}}H_r(X) \longrightarrow \dots$$

For $r > k$, we have $I^{\bar{p}}H_r(\mathcal{U}) = 0$ by Proposition 61. Consider the long exact sequence for relative ordinary homology

$$\dots \longrightarrow H_r(\partial\mathcal{M}) \longrightarrow H_r(\mathcal{M}) \longrightarrow H_r(\mathcal{M}, \partial\mathcal{M}) \longrightarrow \dots$$

We construct a commutative diagram as follows:

(18)

$$\begin{array}{ccccccccc} \dots & \longrightarrow & I^{\bar{p}}H_r(\cap) & \longrightarrow & I^{\bar{p}}H_r(\mathcal{V}) & \longrightarrow & I^{\bar{p}}H_r(X) & \longrightarrow & I^{\bar{p}}H_{r-1}(\cap) & \longrightarrow & I^{\bar{p}}H_{r-1}(\mathcal{V}) & \longrightarrow & \dots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & H_r(\partial\mathcal{M}) & \longrightarrow & H_r(\mathcal{M}) & \longrightarrow & H_r(\mathcal{M}, \partial\mathcal{M}) & \longrightarrow & H_{r-1}(\partial\mathcal{M}) & \longrightarrow & H_{r-1}(\mathcal{M}) & \longrightarrow & \dots \end{array}$$

We look at the morphism $I^{\bar{p}}C_*(\partial\mathcal{M}) \hookrightarrow C_*(\partial\mathcal{M})$ of singular ordinary and intersection chain complexes. Since \mathcal{M} is a \mathbb{Q} -manifold, this morphism induces an isomorphism

$$I^{\bar{p}}H_*(\partial\mathcal{M}) \xrightarrow{\cong} H_*(\partial\mathcal{M}).$$

The outer left vertical homomorphism, which turns out to be an isomorphism, is the composition $I^{\bar{p}}H_*(\cap) \xrightarrow{\cong} I^{\bar{p}}H_*(\partial\mathcal{M}) \xrightarrow{\cong} H_*(\partial\mathcal{M})$. The stratum preserving deformation retraction $\mathcal{V} \rightarrow \mathcal{M}$, induces an isomorphism $I^{\bar{p}}H_*(\mathcal{V}) \xrightarrow{\cong} I^{\bar{p}}H_*(\mathcal{M})$. The morphism $I^{\bar{p}}C_*(\mathcal{M}) \hookrightarrow C_*(\mathcal{M})$ induces an isomorphism $I^{\bar{p}}H_*(\mathcal{M}) \xrightarrow{\cong} H_*(\mathcal{M})$ since \mathcal{M} is a \mathbb{Q} -manifold. The second left vertical homomorphism is the composition $I^{\bar{p}}H_*(\mathcal{V}) \xrightarrow{\cong} I^{\bar{p}}H_*(\mathcal{M}) \xrightarrow{\cong} H_*(\mathcal{M})$. The construction of the two right hand vertical homomorphisms goes along the same lines. Describing the middle

homomorphism requires a bit more effort. First, note that $H_*(\mathcal{M}, \partial\mathcal{M}) \cong \tilde{H}_*(\mathcal{M}/\partial\mathcal{M})$. We show that $H_j(X) \cong H_j(\mathcal{M}/\partial\mathcal{M})$ for $j \geq 2$. We use the Mayer-Vietories sequence for ordinary homology groups. Let $A_{\mathcal{M}/\partial\mathcal{M}} \cong \mathcal{C}(\bigsqcup_i \mathcal{L}_{x_i}) \subset \mathcal{M}/\partial\mathcal{M}$ and $B_{\mathcal{M}/\partial\mathcal{M}} \cong \mathcal{M} \subset \mathcal{M}/\partial\mathcal{M}$ such that $A_{\mathcal{M}/\partial\mathcal{M}} \cap B_{\mathcal{M}/\partial\mathcal{M}} \cong \bigsqcup_i \mathcal{L}_{x_i} \times (0, 1)$. We also set $A_X \cong \bigsqcup_i \mathcal{C}(\mathcal{L}_{x_i}) \subset X$ and $B_X \cong \mathcal{M} \subset X$ such that $A_X \cap B_X \cong \bigsqcup_i \mathcal{L}_{x_i} \times (0, 1)$. For $j \geq 1$, we have

$$H_j(A_X) \oplus H_j(B_X) \cong H_j(A_{\mathcal{M}/\partial\mathcal{M}}) \oplus H_j(B_{\mathcal{M}/\partial\mathcal{M}}) \cong H_j(\mathcal{M}).$$

The claim follows by using 5-lemma. Note that $k \geq 1$. Let $I^{\bar{p}}H_*(X) \rightarrow H_*(X)$ be the homomorphism induced by the morphism $I^{\bar{p}}C_*(X) \hookrightarrow C_*(X)$. The middle vertical homomorphism is the composition

$$I^{\bar{p}}H_*(X) \rightarrow H_*(X) \xrightarrow{\cong} H_*(\mathcal{M}/\partial\mathcal{M}) \xrightarrow{\cong} H_*(\mathcal{M}, \partial\mathcal{M}), \quad * \geq 2.$$

We still need to check the commutativity of the above diagram. Consider the commutative diagram

$$\begin{array}{ccc} \partial\mathcal{M} \times (\frac{1}{2}, 1) & \hookrightarrow & \mathcal{V} \\ \downarrow & & \uparrow \\ \partial\mathcal{M} & \hookrightarrow & \mathcal{M} \end{array} .$$

Each inclusion induces a morphism between intersection chain complexes, and the induced diagram on intersection chain complexes commutes. Hence, the induced diagram on intersection homology groups

$$\begin{array}{ccc} I^{\bar{p}}H_*(\partial\mathcal{M} \times (\frac{1}{2}, 1)) & \longrightarrow & I^{\bar{p}}H_*(\mathcal{V}) \\ \cong \uparrow & & \uparrow \cong \\ I^{\bar{p}}H_*(\partial\mathcal{M}) & \longrightarrow & I^{\bar{p}}H_*(\mathcal{M}) \end{array}$$

commutes. Recall that vertical homomorphisms are isomorphisms. Furthermore, the diagram

$$\begin{array}{ccc} I^{\bar{p}}H_*(\partial\mathcal{M}) & \longrightarrow & I^{\bar{p}}H_*(\mathcal{M}) \\ \cong \downarrow & & \downarrow \cong \\ H_*(\partial\mathcal{M}) & \longrightarrow & H_*(\mathcal{M}) \end{array}$$

commutes. Note that the vertical morphisms induced by inclusions of singular intersection chain complexes into ordinary singular chain complexes are isomorphisms, as mentioned. Hence, the outer left square commutes. Using the same procedure, one can show that the outer right square commutes. Consider the commutative diagram

$$\begin{array}{ccc} \mathcal{V} & \hookrightarrow & X \\ \uparrow & \nearrow & \\ \mathcal{M} & & \end{array} .$$

Each inclusion induces a morphism between intersection chain complexes, and the induced diagram on intersection chain complexes commutes. Hence, the induced diagram on intersection homology groups

$$\begin{array}{ccc} I^{\bar{p}}H_*(\mathcal{V}) & \longrightarrow & I^{\bar{p}}H_*(X) \\ \uparrow & \nearrow & \\ I^{\bar{p}}H_*(\mathcal{M}) & & \end{array}$$

commutes. The commutativity of

$$\begin{array}{ccc} I^{\bar{p}}C_*(\mathcal{M}) & \longrightarrow & I^{\bar{p}}C_*(X) \\ \downarrow & & \downarrow \\ C_*(\mathcal{M}) & \longrightarrow & C_*(X) \end{array}$$

ensures that

$$\begin{array}{ccc} I^{\bar{p}}H_*(\mathcal{M}) & \longrightarrow & I^{\bar{p}}H_*(X) \\ \downarrow & & \downarrow \\ H_*(\mathcal{M}) & \longrightarrow & H_*(X) \end{array}$$

commutes. Thus, the second left square commutes. Consider the short exact sequence

$$0 \longrightarrow I^{\bar{p}}C_i(\cap) \longrightarrow I^{\bar{p}}C_i(\mathcal{U}) \oplus I^{\bar{p}}C_i(\mathcal{V}) \longrightarrow I^{\bar{p}}C_i(\mathcal{U}) + I^{\bar{p}}C_i(\mathcal{V}) \longrightarrow 0.$$

Note that $I^{\bar{p}}C_i(\mathcal{U}) + I^{\bar{p}}C_i(\mathcal{V})$ is the sub-complex of $I^{\bar{p}}C_i(X)$ generated by allowable chains supported in \mathcal{U} or in \mathcal{V} . The commutativity of the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & I^{\bar{p}}C_i(\cap) & \longrightarrow & I^{\bar{p}}C_i(\mathcal{U}) \oplus I^{\bar{p}}C_i(\mathcal{V}) & \longrightarrow & I^{\bar{p}}C_i(\mathcal{U}) + I^{\bar{p}}C_i(\mathcal{V}) \longrightarrow 0. \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_i(\cap) & \longrightarrow & C_i(\mathcal{U}) \oplus C_i(\mathcal{V}) & \longrightarrow & C_i(\mathcal{U}) + C_i(\mathcal{V}) \longrightarrow 0 \end{array}$$

implies that the diagram

$$\begin{array}{ccccccc} \dots & \longrightarrow & I^{\bar{p}}H_r(\mathcal{V}) & \longrightarrow & I^{\bar{p}}H_r(X) & \longrightarrow & I^{\bar{p}}H_{r-1}(\cap) \longrightarrow \dots \\ & & \downarrow & & \downarrow & & \downarrow \\ \dots & \longrightarrow & H_r(\mathcal{V}) & \longrightarrow & H_r(X) & \longrightarrow & H_{r-1}(\cap) \longrightarrow \dots \end{array}$$

commutes. The commutativity of the right square in the above diagram ensures that the second right square in Diagram 18 commutes. The desired result follows from 5-lemma. Thus, we have

$$I^{\bar{p}}H_r(X) \cong H_r(\mathcal{M}, \partial\mathcal{M}), \text{ for } r > k.$$

For $r < k$, the proof goes as follows. Let \mathcal{U} and \mathcal{V} be as above. Employing Proposition 61 yields $I^{\bar{p}}H_r(\mathcal{U}) \cong I^{\bar{p}}H_r(\mathcal{L})$. Recall that \mathcal{L} is a \mathbb{Q} -manifold. Hence, we have the following diagram.

$$\begin{array}{ccccccccccc} \dots & \rightarrow & I^{\bar{p}}H_r(\cap) & \rightarrow & I^{\bar{p}}H_r(\mathcal{M}) \oplus I^{\bar{p}}H_r(\mathcal{L}) & \rightarrow & I^{\bar{p}}H_r(X) & \rightarrow & I^{\bar{p}}H_{r-1}(\cap) & \rightarrow & I^{\bar{p}}H_{r-1}(\mathcal{M}) \oplus I^{\bar{p}}H_{r-1}(\mathcal{L}) & \rightarrow \dots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\ \dots & \rightarrow & H_r(\cap) & \rightarrow & H_r(\mathcal{M}) \oplus H_r(\mathcal{L}) & \rightarrow & H_r(\mathcal{M}) & \rightarrow & H_{r-1}(\cap) & \rightarrow & H_{r-1}(\mathcal{M}) \oplus H_{r-1}(\mathcal{L}) & \rightarrow \dots \end{array}$$

The commutativity of the left hand and right hand squares follows from the commutativity of

$$\begin{array}{ccc} I^{\bar{p}}C_*(\cap) & \longrightarrow & I^{\bar{p}}C_*(\mathcal{L}) \oplus I^{\bar{p}}C_*(\mathcal{M}) . \\ \downarrow & & \downarrow \\ C_*(\cap) & \longrightarrow & C_*(\mathcal{L}) \oplus C_*(\mathcal{M}) \end{array}$$

The vertical monomorphisms induce isomorphisms between the intersection and ordinary homology groups. By [6, Lemma 2.46], we can construct the middle vertical homomorphism such that the diagram commutes. Finally, by 5-lemma, we have

$$I^{\bar{p}}H_r(X) \cong H_r(\mathcal{M}), \text{ for } r < k.$$

This concludes the proof. \square

We are now at the point where we can start with the proof of the main theorem of this section. In [6], the first named author shows more than just the duality of Betti numbers of intersection spaces. Although we are only interested in the duality of the Betti numbers of intersection spaces, we can generalize [6, Theorem 2.12] to \mathbb{Q} -pseudomanifolds. For a brief introduction to reflective algebra, the reader may consult [6, Section 2.1].

Remark 63. *As mentioned earlier, we only consider homology with rational coefficients. Note also that because the underlying space, X , is a \mathbb{Q} -pseudomanifold with \mathbb{Q} -isolated singularities, $I^{\bar{p}}X$ refers to the generalized intersection space of X in the sense of Definition 60.*

Theorem 64. *Let X be an n -dimensional compact oriented \mathbb{Q} -pseudomanifold with only \mathbb{Q} -isolated singularities. Let \bar{p} and \bar{q} be complementary perversities. Assume that the link of each \mathbb{Q} -isolated singularity is rationally $(n - 1 - \bar{p}(n))$ -segmented.*

Then

- (1) *The pair $(\tilde{H}_*(I^{\bar{p}}X), I^{\bar{p}}H_*(X))$ is $(n - 1 - \bar{p}(n))$ -reflective across the homology of the links and*
- (2) *$(\tilde{H}_*(I^{\bar{p}}X), I^{\bar{p}}H_*(X))$ and $(\tilde{H}_*(I^{\bar{q}}X), I^{\bar{q}}H_*(X))$ are n -dual reflective pairs.*

Proof. We mainly mimic the proof of [6, Theorem 2.12]. By the above considerations, the main ingredients of the proof remain intact for \mathbb{Q} -Pseudomanifolds with \mathbb{Q} -isolated singularities.

We start with the homology braid of the triple

$$\mathcal{L}_{<k} \hookrightarrow \mathcal{L} \xrightarrow{j} \mathcal{M}.$$

By Proposition 62, the same argument as in the proof of [6, Theorem 2.12] can be applied. The rest of the proof relies on Lefschetz duality for $(\mathcal{M}, \partial\mathcal{M})$ and Poincaré duality for $\partial\mathcal{M}$. For Lefschetz duality, we employ Theorem 47, The topological space $\partial\mathcal{M}$ is a \mathbb{Q} -manifold by Proposition 50, and therefore it satisfies Poincaré duality rationally, using Corollary 49. This concludes our proof. \square

Corollary 65. *Let X be an n -dimensional, compact, oriented, topological \mathbb{Q} -pseudomanifold with only \mathbb{Q} -isolated singularities, which means there is a \mathbb{Q} -stratification of the form $X_n \supset X_0$. Let \bar{p} and \bar{q} be complementary perversities. Then, we have the duality isomorphism*

$$d : \tilde{H}_r(I^{\bar{p}}X)^* \xrightarrow{\cong} \tilde{H}_{n-r}(I^{\bar{q}}X),$$

where

$$\tilde{H}_r(I^{\bar{p}}X)^* = \text{Hom}(\tilde{H}_r(I^{\bar{p}}X), \mathbb{Q}).$$

Proof. The construction of the duality isomorphism goes along the same line as in the proof of [6, Theorem 2.12] and using Corollary 49 and Theorem 47. \square

Corollary 58 continues to hold in the \mathbb{Q} -isolated setting for the generalized intersection spaces:

Corollary 66. *Let X be an n -dimensional, compact, oriented \mathbb{Q} -pseudomanifold with only \mathbb{Q} -isolated singularities. Let $I^{\bar{p}}X$ be the associated generalized intersection spaces. If $n = \dim(X)$ is even, then the difference between the Euler characteristic of $\tilde{H}_*(I^{\bar{p}}X)$ and $I^{\bar{p}}H_*(X)$ is given by*

$$\chi(\tilde{H}_*(I^{\bar{p}}X)) - \chi(I^{\bar{p}}H_*(X)) = -2\chi_{<n-1-\bar{p}(n)}(\mathcal{L}),$$

where \mathcal{L} is the disjoint union of the links of the \mathbb{Q} -isolated singularities of X .
If $n = \dim(X)$ is odd, then

$$\chi(\widetilde{H}_*(I^{\bar{n}}X)) - \chi(I^{\bar{n}}H_*(X)) = (-1)^{\frac{n-1}{2}} b_{\frac{n-1}{2}}(\mathcal{L}),$$

where $b_{\frac{n-1}{2}}(\mathcal{L})$ is the middle dimensional Betti number of \mathcal{L} and \bar{n} is the upper middle perversity.
Regardless of the parity of n , the identity

$$\mathrm{rk}(\widetilde{H}_k(I^{\bar{p}}X)) + \mathrm{rk}(I^{\bar{p}}H_k(X)) = \mathrm{rk}(H_k(\mathcal{M})) + \mathrm{rk}(H_k(\mathcal{M}, \mathcal{L}))$$

always holds in degree $k = n - 1 - \bar{p}(n)$, where \mathcal{M} is the \mathbb{Q} -manifold obtained by cutting out the \mathbb{Q} -isolated singularities of X .

8. INTERSECTION SPACES OF 6-DIMENSIONAL TORIC VARIETIES

As mentioned in Section 4.3, a 6-dimensional toric variety is a \mathbb{Q} -pseudomanifold with \mathbb{Q} -isolated singularities. Such varieties are therefore amenable to the theory developed in Section 7. Although we will not compute the homology groups of the intersection spaces directly via CW structures, we will show that considering the middle perversity, links of \mathbb{Q} -isolated singularities are rationally 3-segmented. Using Lefschetz duality, we then compute the rational homology groups of generalized intersection spaces of 6-dimensional toric varieties. As a side remark, we will also show that Corollary 66 yields the well-known Euler formula for 3-dimensional polytopes. Let $X_{\mathcal{P}}$ be a 6-dimensional toric variety associated with \mathcal{P} , the underlying polytope. We can endow $X_{\mathcal{P}}$ with a \mathbb{Q} -stratification of the form $X_{\mathcal{P}} = X_6 \supset X_0$. We have shown in Section 4.3 that links of \mathbb{Q} -isolated singularities are \mathbb{Q} -manifolds and satisfy Poincaré duality rationally.

Proposition 67. *Let X be a 6-dimensional toric variety with a \mathbb{Q} -stratification of the form $X_6 \supset X_0$. For each $x_i \in X_0$, let \mathcal{L}_i be the associated link. Then \mathcal{L}_i is rationally 3-segmented.*

Proof. We endow each \mathcal{L}_i with the CW structure described in Section 4.3. Recall that we have computed ∂_3 of the \mathcal{L}_i in Equation 13. From each T^2 with one 2-cell, we remove this 2-cell. From each T^2 with two 2-cells, we omit the 2-cell with the negative sign in the boundary operator. We denote the obtained 3-dimensional CW sub-complex by \mathcal{L}'_i . Despite the removal of the 2-cells from the tori, the 2-skeleton of the link has not changed, i.e \mathcal{L}_i and \mathcal{L}'_i have the same 2-skeleton. Hence, we get

$$(19) \quad \partial_3^{\mathcal{L}'_i} = \begin{matrix} e_{T_3}^0 \times e_{\mathcal{M}_x}^2 \\ e_{(T_{a_i}^2)_1}^1 \times e_{(\mathcal{M}_{x_a})_i}^1 \tilde{a} \\ e_{(T_{a_i}^2)_2}^1 \times e_{(\mathcal{M}_{x_a})_i}^1 \tilde{a} \\ e_{(T_{a_i}^2)_3}^1 \times e_{(\mathcal{M}_{x_a})_i}^1 \tilde{a} \\ e_{(T_{b_j}^2)_1}^1 \times e_{(\mathcal{M}_{x_b})_j}^1 \tilde{b} \\ e_{(T_{b_j}^2)_2}^1 \times e_{(\mathcal{M}_{x_b})_j}^1 \tilde{b} \end{matrix} \begin{bmatrix} e_{T_3}^1 \times e_{\mathcal{M}_x}^2 & e_{T_2}^1 \times e_{\mathcal{M}_x}^2 & e_{T_3}^1 \times e_{\mathcal{M}_x}^2 & e_{(T_{a_i}^2)_1}^2 \times e_{(\mathcal{M}_{x_a})_i}^1 \\ 0 & 0 & 0 & \overbrace{0}^{\tilde{a}} \\ m_{a_i} l_{a_i} & 0 & 0 & 1 \\ 0 & n_{a_i} l_{a_i} & 0 & 1 \\ 0 & 0 & n_{a_i} m_{a_i} & 1 \\ -m_{b_j} & 0 & 0 & 0 \\ 0 & n_{b_j} & 0 & 0 \end{bmatrix}.$$

We use the same notation as in Section 4.3. Recall that each 1-dimensional face of \mathcal{M}_x , is associated to a 2-dimensional face of \mathcal{P} , which is dual to a 1-dimensional cone in Σ such that the dual cones lie in \mathcal{S}_x . Here \tilde{a} and \tilde{b} are the numbers of such 1-dimensional cones in Σ , whose generators have no zero entry and at least one zero-entry, respectively. Recall also that $e_{(\mathcal{M}_{x_a})_i}^1$ and $e_{(\mathcal{M}_{x_b})_j}^1$ are those 1-cells of \mathcal{M}_x , whose associated T^2 in \mathcal{L}_x has three and two 1-cells,

respectively. Clearly, we have

$$\mathrm{Im}(\partial_3^{\mathcal{L}'_i}) = \mathrm{Im}(\partial_3^{\mathcal{L}_i}) \text{ and } \ker(\partial_3^{\mathcal{L}'_i}) = 0,$$

and the inclusion $\mathrm{incl} : \mathcal{L}'_i \hookrightarrow \mathcal{L}_i$ induces

$$\mathrm{incl}_j : H_j(\mathcal{L}'_i) \xrightarrow{\cong} H_j(\mathcal{L}_i) \text{ for } j \leq 2.$$

Obviously,

$$H_j(\mathcal{L}'_i) = 0 \text{ for } j > 2.$$

□

Theorem 68. *Let $X_{\mathcal{P}}$ be an arbitrary compact 6-dimensional toric variety with m \mathbb{Q} -isolated singularities, $m \geq 1$, where \mathcal{P} is the underlying polytope. Let Σ be the dual fan to \mathcal{P} . We denote the number of 1-dimensional and 2-dimensional cones of Σ with f_1 and f_2 , respectively. Then,*

$$\begin{aligned} \mathrm{rk}(\tilde{H}_6(I^{\bar{n}}X)) &= 0 \\ \mathrm{rk}(\tilde{H}_5(I^{\bar{n}}X)) &= m - 1 \\ \mathrm{rk}(\tilde{H}_4(I^{\bar{n}}X)) &= f_1 - 3 - b \\ \mathrm{rk}(\tilde{H}_3(I^{\bar{n}}X)) &= 2(3f_1 - f_2 - b - 6) \\ \mathrm{rk}(\tilde{H}_2(I^{\bar{n}}X)) &= f_1 - 3 - b \\ \mathrm{rk}(\tilde{H}_1(I^{\bar{n}}X)) &= m - 1 \\ \mathrm{rk}(\tilde{H}_0(I^{\bar{n}}X)) &= 0, \end{aligned}$$

where $I^{\bar{n}}X$ is the generalized intersection space associated to the \mathbb{Q} -pseudomanifold $X_{\mathcal{P}}$, and the parameter b is introduced and has been studied in Section 4.2.1.

Proof. We start by cutting out the \mathbb{Q} -isolated singularities of $X_{\mathcal{P}}$, in the sense of Definition 41. We denote the resulting \mathbb{Q} -manifold with boundary by $(\mathcal{M}, \partial\mathcal{M})$. It satisfies Lefschetz duality rationally by Corollary 48. Now, let $\mathcal{M}/\partial\mathcal{M}$ be the topological pseudomanifold obtained by coning off the boundary of $(\mathcal{M}, \partial\mathcal{M})$. As shown, we have

$$H_i(X) \cong H_i(\mathcal{M}/\partial\mathcal{M}) \cong H_i(\mathcal{M}, \partial\mathcal{M}) \text{ for } i \geq 2.$$

Hence by Proposition 27, we get

$$\begin{aligned} \mathrm{rk}(H_6(\mathcal{M}, \partial\mathcal{M})) &= \mathrm{rk}(H_0(\mathcal{M})) = 1 \\ \mathrm{rk}(H_5(\mathcal{M}, \partial\mathcal{M})) &= \mathrm{rk}(H_1(\mathcal{M})) = 0 \\ \mathrm{rk}(H_4(\mathcal{M}, \partial\mathcal{M})) &= \mathrm{rk}(H_2(\mathcal{M})) = f_1 - 3 \\ \mathrm{rk}(H_3(\mathcal{M}, \partial\mathcal{M})) &= \mathrm{rk}(H_3(\mathcal{M})) = 3f_1 - f_2 - b - 6 \\ \mathrm{rk}(H_2(\mathcal{M}, \partial\mathcal{M})) &= \mathrm{rk}(H_4(\mathcal{M})) = f_1 - 3 - b. \end{aligned}$$

The group $H_6(\mathcal{M})$ vanishes since by Lefschetz duality $\mathrm{rk}(H_6(\mathcal{M})) = \mathrm{rk}(H_0(\mathcal{M}, \partial\mathcal{M})) = 0$ (Note that $\partial\mathcal{M}$ is not empty as $m \geq 1$.) Consequently, by considering the long exact sequence of the pair $(\mathcal{M}, \partial\mathcal{M})$, we arrive at the short exact sequence

$$0 \longrightarrow H_6(\mathcal{M}, \partial\mathcal{M}) \longrightarrow H_5(\partial\mathcal{M}) \longrightarrow H_5(\mathcal{M}) \longrightarrow 0.$$

Since $\partial\mathcal{M} = \bigsqcup_{i=1}^m \mathcal{L}_i$ and each \mathcal{L}_i is orientable, connected and 5-dimensional, $\mathrm{rk}(H_5(\partial\mathcal{M})) = m$. This implies

$$\mathrm{rk}(H_1(\mathcal{M}, \partial\mathcal{M})) = \mathrm{rk}(H_5(\mathcal{M})) = m - 1.$$

Before dealing with the homology groups of the generalized intersection space, we need to compute the intersection homology groups of $X_{\mathcal{P}}$. In [32], Stanley shows that the odd degree

intersection homology groups of toric varieties vanish. We merely need this result for the middle degree. The relevant cut off degree is $k = 6 - 1 - \bar{n}(6) = 3$. Using Proposition 62, we finally get

$$\begin{aligned} \operatorname{rk}(I^{\bar{n}}H_6(X)) &= 1 \\ \operatorname{rk}(I^{\bar{n}}H_5(X)) &= 0 \\ \operatorname{rk}(I^{\bar{n}}H_4(X)) &= f_1 - 3 \\ \operatorname{rk}(I^{\bar{n}}H_3(X)) &= 0 \\ \operatorname{rk}(I^{\bar{n}}H_2(X)) &= f_1 - 3 \\ \operatorname{rk}(I^{\bar{n}}H_1(X)) &= 0 \\ \operatorname{rk}(I^{\bar{n}}H_0(X)) &= 1. \end{aligned}$$

Proposition 67 ensures the existence of the associated generalized intersection space of $X_{\mathcal{P}}$. By using the duality isomorphisms of Theorem 64, we can read off the homology groups of $I^{\bar{n}}X$ above and below the middle degree:

$$\begin{aligned} H_r(\mathcal{M}) &\xrightarrow{\cong} \tilde{H}_r(I^{\bar{n}}X) \text{ for } r > k = 3 \\ H_r(\mathcal{M}, \partial\mathcal{M}) &\xrightarrow{\cong} \tilde{H}_r(I^{\bar{n}}X) \text{ for } r < k = 3. \end{aligned}$$

Finally, we use Proposition 33 and the last identity in Corollary 66 and compute $\operatorname{rk}(\tilde{H}_3(I^{\bar{n}}X))$. \square

Corollary 69. *Let $X_{\mathcal{P}}$ be an arbitrary compact 6-dimensional toric variety, where \mathcal{P} is the underlying polytope. Let Σ be the fan dual to \mathcal{P} . We denote the 1-dimensional, 2-dimensional, and 3-dimensional cones of Σ with f_1 , f_2 , and f_3 , respectively. We further assume that all vertices in \mathcal{P} are \mathbb{Q} -isolated singularities. Then,*

$$\begin{aligned} \operatorname{rk}(\tilde{H}_6(I^{\bar{n}}X)) &= 0 \\ \operatorname{rk}(\tilde{H}_5(I^{\bar{n}}X)) &= f_3 - 1 \\ \operatorname{rk}(\tilde{H}_4(I^{\bar{n}}X)) &= f_1 - 3 - b \\ \operatorname{rk}(\tilde{H}_3(I^{\bar{n}}X)) &= 2(3f_1 - f_2 - b - 6) \\ \operatorname{rk}(\tilde{H}_2(I^{\bar{n}}X)) &= f_1 - 3 - b \\ \operatorname{rk}(\tilde{H}_1(I^{\bar{n}}X)) &= f_3 - 1 \\ \operatorname{rk}(\tilde{H}_0(I^{\bar{n}}X)) &= 0. \end{aligned}$$

Proof. The number of vertices in \mathcal{P} is equal to f_3 , which equals m by the assumption. \square

Remark 70. *For $X_{\mathcal{P}}$ an arbitrary compact 6-dimensional toric variety with m \mathbb{Q} -isolated singularities, $m \geq 1$, we employ the first identity on the Euler characteristic in Corollary 66 to obtain*

$$\begin{aligned} &(0 - (m - 1) + (f_1 - 3 - b) - 2(3f_1 - f_2 - b - 6) + (f_1 - 3 - b) \\ &- (m - 1) + 0) - (1 + (f_1 - 3) + (f_1 - 3) + 1) \\ &= -2\left(\sum_{i=1}^m 1 - 0 + \sum_{i=1}^m (f_1^i - 3)\right), \end{aligned}$$

where f_1^i denotes the number of 1-dimensional neighboring faces of the i -th vertex of \mathcal{P} . We can also interpret f_1^i as the number of 1-dimensional faces of \mathcal{M}_i , where \mathcal{M}_i is the underlying 2-dimensional polygon of the link of the i -th vertex. Note that for a \mathbb{Q} -isolated non-singular point

x_j , we have $f_1^j = 3$. This implies $\sum_{i=1}^m (f_1^i - 3) = \sum_{i=1}^{f_3} (f_1^i - 3)$. Counting the 1-dimensional faces of \mathcal{M}_i for each vertex x_i equals counting the neighboring 1-dimensional faces of each vertex in the underlying polytope \mathcal{P} . Since each 1-dimensional face of the polytope has precisely two vertices in its topological boundary, we have $\sum_{i=1}^{f_3} f_1^i = 2f_2$. Thus, we get

$$\sum_{i=1}^{f_3} (f_1^i - 3) = 2f_2 - 3f_3.$$

Then, we have $(-2m - 4f_1 + 2f_2 + 8) - (2f_1 - 4) = -2(m + 2f_2 - 3f_3)$. Finally, we get

$$f_3 - f_2 + f_1 = 2,$$

which is the well-known Euler formula for 3-dimensional polytopes.

Corollary 71. *We assume that all vertices in \mathcal{P} are \mathbb{Q} -isolated singularities. Considering the above remark, we arrive at*

$$\begin{aligned} \text{rk}(\tilde{H}_6(I^{\bar{n}}X)) &= 0 \\ \text{rk}(\tilde{H}_5(I^{\bar{n}}X)) &= f_2 - f_1 + 1 \\ \text{rk}(\tilde{H}_4(I^{\bar{n}}X)) &= f_1 - 3 - b \\ \text{rk}(\tilde{H}_3(I^{\bar{n}}X)) &= 2(3f_1 - f_2 - b - 6) \\ \text{rk}(\tilde{H}_2(I^{\bar{n}}X)) &= f_1 - 3 - b \\ \text{rk}(\tilde{H}_1(I^{\bar{n}}X)) &= f_2 - f_1 + 1 \\ \text{rk}(\tilde{H}_0(I^{\bar{n}}X)) &= 0. \end{aligned}$$

Remark 72. *In Theorem 68, we have*

$$\chi(\tilde{H}_*(I^{\bar{n}}X)) = 2(-2f_1 + f_2 - m + 4),$$

which is even. This observation is consistent with the following more general principle. Let X be a $(4n+2)$ -dimensional \mathbb{Q} -pseudomanifold with \mathbb{Q} -isolated singularities. By using the duality on the intersection homology groups, we find

$$\chi(I^{\bar{n}}H_*(X)) = \sum_{i=1}^{4n+2} (-1)^i \text{rk}(I^{\bar{n}}H_i(X)) = 2 \sum_{i=1}^{2n} (-1)^i \text{rk}(I^{\bar{n}}H_i(X)) - \text{rk}(I^{\bar{n}}H_{2n+1}(X)).$$

The intersection form on $I^{\bar{n}}H_{2n+1}(X)$ is skew-symmetric and non-degenerate. Hence, there is a basis with respect to which the intersection form has the matrix representation

$$\begin{pmatrix} 0 & 1 & & & & 0 \\ -1 & 0 & & & & \\ & & 0 & 1 & & \vdots \\ & & -1 & 0 & & \\ \vdots & & & & \ddots & 0 \\ 0 & & \cdots & & 0 & 1 \\ & & & & -1 & 0 \end{pmatrix}.$$

Consequently, we have $\text{rk}(I^{\bar{n}}H_{2n+1}(X)) \equiv 0(2)$ and thus $\chi(I^{\bar{n}}H_*(X)) \equiv 0(2)$. For an intersection space $I^{\bar{n}}X$ of X , it follows from Lefschetz duality that

$$\text{rk}(H_{2n+1}(I^{\bar{n}}X)) = -\text{rk}(I^{\bar{n}}H_{2n+1}(X)) + 2 \text{rk}(H_{2n+1}(\mathcal{M})).$$

Therefore,

$$\chi(H_*(I^{\bar{n}}X)) \equiv 0(2).$$

Remark 73. *In the context of Corollary 71, the homology groups of $I^{\bar{n}}X$ determine the ordinary homology groups of $X_{\mathcal{P}}$ up to and including the middle degree. In contrast, the intersection homology groups do not determine the ordinary homology below the middle degree. In this sense, the homology of intersection spaces of toric varieties sees more structure than the intersection homology.*

Remark 74. *We can nicely fit all of the above data into a Mayer-Vietoris sequence. Let $I^{\bar{n}}X$ be the associated generalized intersection space of a 6-dimensional toric variety X . Then we take $\mathcal{U} \simeq \mathcal{M}$, where \mathcal{M} is the \mathbb{Q} -manifold obtained by cutting out the \mathbb{Q} -isolated singularities of X . We also take $\mathcal{V} \cong \mathcal{C}(\sqcup \mathcal{L}_{<3})$. The intersection $\cap := \mathcal{U} \cap \mathcal{V}$ is homotopy equivalent to $\mathcal{L}_{<3}$. Thus, we have*

$$\begin{array}{ccccccccccc}
0 & \longrightarrow & \overbrace{H_6(\cap)}^{=0} & \longrightarrow & \overbrace{H_6(\mathcal{U})}^{=0} & \xrightarrow{\cong} & \overbrace{H_6(I^{\bar{n}}X)}^{=0} & \longrightarrow & \overbrace{H_5(\cap)}^{=0} & \longrightarrow & \overbrace{H_5(\mathcal{U})}^{\cong \mathbb{Q}^{m-1}} & \xrightarrow{\cong} & \overbrace{H_5(I^{\bar{n}}X)}^{\cong \mathbb{Q}^{m-1}} \\
& & \overbrace{H_4(\cap)}^{=0} & \longrightarrow & \overbrace{H_4(\mathcal{U})}^{\cong \mathbb{Q}^{f_1-3-b}} & \xrightarrow{\cong} & \overbrace{H_4(I^{\bar{n}}X)}^{\cong \mathbb{Q}^{f_1-3-b}} & \longrightarrow & \overbrace{H_3(\cap)}^{=0} & \longrightarrow & \overbrace{H_3(\mathcal{U})}^{\cong \mathbb{Q}^{3f_1-f_2-b-6}} & \longrightarrow & \overbrace{H_3(I^{\bar{n}}X)}^{\cong \mathbb{Q}^{2(3f_1-f_2-b-6)}} \\
& & \overbrace{H_2(\cap)}^{\cong \mathbb{Q}^r} & \longrightarrow & \overbrace{H_2(\mathcal{U})}^{\cong \mathbb{Q}^{f_1-3}} & \longrightarrow & \overbrace{H_2(I^{\bar{n}}X)}^{\cong \mathbb{Q}^{f_1-3-b}} & \longrightarrow & \overbrace{H_1(\cap)}^{=0} & \longrightarrow & \overbrace{H_1(\mathcal{U})}^{=0} & \longrightarrow & \overbrace{H_1(I^{\bar{n}}X)}^{\cong \mathbb{Q}^{m-1}} \\
& & \overbrace{H_0(\cap)}^{\cong \mathbb{Q}^{m-1}} & \xrightarrow{\cong} & \overbrace{H_0(\mathcal{U})}^{=0} \oplus \overbrace{H_0(\mathcal{V})}^{=0} & \longrightarrow & \overbrace{H_0(I^{\bar{n}}X)}^{=0} & \longrightarrow & 0, & & & &
\end{array}$$

where $r := \sum_{i=1}^m (f_1^i - 3) = 2f_2 - 3f_3$.

Corollary 75. *For each 6-dimensional toric variety X with $H_3(X) = 0$, the equality $b = r$ holds and b is combinatorially invariant.*

Proof. The assumption implies that $H_3(\mathcal{U}) = 0$ and hence $H_3(I^{\bar{n}}X) = 0$. By using the Mayer-Vietoris sequence of Remark 74, we get $b = r = 2f_2 - 3f_3$. Reformulating the assumption yields $b = 3f_1 - f_2 - 6$. Setting both expressions equal to each other will result in the Euler formula. \square

9. CONCLUSION

We summarize the results obtained in this work. We started by considering toric varieties as topological pseudomanifolds. We introduced the construction of a link for an arbitrary stratum in a given toric variety. We also endowed links of strata with real co-dimensions 4 and 6 with CW structures and computed their homology groups. We showed that the link of each connected component of the real 4-co-dimensional stratum is always a rational homology 3-sphere. Thus, for an arbitrary $2n$ -dimensional toric variety X , we get a \mathbb{Q} -stratification whose depth 1 stratum is of real co-dimension 6. We generalized the theory of intersection spaces to rational homology stratified pseudomanifolds with rationally isolated singularities. Our main objects of study are 6-dimensional toric varieties. For these, we compare the Betti numbers of the associated generalized intersection spaces IX and intersection homology in the following table.

The left hand column of the table shows that the b_*^{IX} are not combinatorially invariant. Indeed, we can determine the ordinary Betti numbers of X up to and including the middle degree by using b_*^{IX} . However, the right hand column, as already observed by Fieseler in [18], shows that the associated Betti numbers of intersection homology neither fix the combinatorial data of the fan nor the ordinary homology below the middle degree.

b_* \ Theory	$H_*(IX)$	$IH_*(X)$
b_6	0	1
b_5	$m - 1$	0
b_4	$f_1 - 3 - b$	$f_1 - 3$
b_3	$2(3f_1 - f_2 - b - 6)$	0
b_2	$f_1 - 3 - b$	$f_1 - 3$
b_1	$m - 1$	0
b_0	1	1

TABLE 1. The Betti numbers associated to HI_* and IH_* .

REFERENCES

- [1] Agustín, M., Essig, J. & de Bobadilla, J. *Self-dual intersection space complexes*, (2020) [arXiv: 2005.14616](#)
- [2] Agustín Vicente, M. & de Bobadilla, J. *Intersection space constructible complexes*, Documenta Mathematica, **25** (2020), pp. 1653–1725 [DOI: 10.4171/dm/783](#)
- [3] Banagl, M. *Extending Intersection Homology Type Invariants to Non-Witt Spaces*, Memoirs of the Amer. Math. Soc. **160**, no. 760, American Math. Society, 2002. [DOI: 10.1090/memo/0760](#)
- [4] Banagl, M. *Topological invariants of stratified spaces*, Springer Monographs in Mathematics, Springer-Verlag Berlin Heidelberg (2007)
- [5] Banagl, M. *Foliated stratified spaces and a de Rham complex describing intersection space cohomology*, Journal Of Differential Geometry, **104** (2016), pp. 1–58 [DOI: 10.4310/jdg/1473186538](#)
- [6] Banagl, M. *Intersection spaces, spatial homology truncation, and string theory*, Lecture Notes in Mathematics **1997**. (Springer-Verlag Berlin - Heidelberg, 2010) [DOI: 10.1007/978-3-642-12589-8](#)
- [7] Banagl, M. *Isometric group actions and the cohomology of flat fiber bundles*, Groups Geometry And Dynamics, **7** (2013), pp. 293–321 [DOI: 10.4171/ggd/183](#)
- [8] Banagl, M., Budur, N. & Maxim, L. *Intersection spaces, perverse sheaves and type IIB string theory*, Advances In Theoretical And Mathematical Physics, **18** (2014), pp. 363-399 [DOI: 10.4310/ATMP.2014.v18.n2.a3](#)
- [9] Banagl, M. & Hunsicker, E. *Hodge theory for intersection space cohomology*, Geometry & Topology, **23** (2019), pp 2165–2225 [DOI: 10.2140/gt.2019.23.2165](#)
- [10] Banagl, M. & Maxim, L. *Intersection spaces and hypersurface singularities*, Journal Of Singularities, **5** (2012), pp. 48–56 [DOI: 10.5427/jsing.2012.5d](#)
- [11] Borel, A., et al., *Intersection Cohomology*, Progress in Math., vol. 50, Birkhäuser, 1984. [DOI: 10.1007/978-0-8176-4765-0](#)
- [12] Cheeger, J. *On the Hodge theory of Riemannian pseudomanifolds*, in Proc. of Symposia in Pure Math, **36** (1980), pp. 91–146, American Math. Soc. [DOI: 10.1090/pspum/036/573430](#)
- [13] Cheeger, J. *On the spectral geometry of spaces with cone-like singularities*, Proceedings Of The National Academy Of Sciences, **76** (1979), pp. 2103–2106 [DOI: 10.1073/pnas.76.5.2103](#)
- [14] Cheeger, J. *Spectral geometry of singular Riemannian spaces*, Journal Of Differential Geometry, **18** (1983), pp. 575–657 [DOI: 10.4310/jdg/1214438175](#)
- [15] Cox, D., Little, J. & Schenck, H. *Toric varieties*, (American Mathematical Soc., 2011) [DOI: 10.1090/gsm/124](#)
- [16] Essig, T. *About a de Rham Complex describing intersection space cohomology in a three strata case*, PhD thesis, (Universität Heidelberg 2016)
- [17] Essig, T. *A generalized de Rham Theorem for Intersection Spaces*, Matemática Contemporânea, **53** (2023), pp. 233–253 [DOI: 10.21711/231766362023/rmc5311](#)
- [18] Fieseler, K. *Rational intersection cohomology of projective toric varieties*, J. für die reine und angewandte Mathematik, **413** (1991), pp. 88–98 [DOI: 10.1515/crll.1991.413.88](#)
- [19] Fischli, S. *On toric varieties*. PhD thesis. (Universität Bern, 1992)
- [20] Friedman, G. *Singular intersection homology*. (Cambridge University Press, 2020) [DOI: 10.1017/9781316584446](#)
- [21] Friedman, G. & McClure, J. *Cup and cap products in intersection (co)homology*, Advances In Mathematics, **240** (2013), pp. 383–426 [DOI: 10.1016/j.aim.2013.02.017](#)
- [22] Goresky, M. & MacPherson, R. *Intersection homology theory*, Topology, **19** (1980), pp. 135–162 [DOI: 10.1016/0040-9383\(80\)90003-8](#)

- [23] Goresky, M. & MacPherson, R. *Intersection Homology II*, Invent. Math. **71** (1983), 77 – 129. DOI: [10.1007/BF01389130](https://doi.org/10.1007/BF01389130)
- [24] Iversen, B. *Cohomology of Sheaves*, Universitext, Springer-Verlag Berlin Heidelberg 1986. DOI: [10.1007/978-3-642-82783-9](https://doi.org/10.1007/978-3-642-82783-9)
- [25] Kirwan, F. & Wolf, J. *An introduction to intersection homology theory*, (CRC Press, 2006) DOI: [10.1201/9780367800840](https://doi.org/10.1201/9780367800840)
- [26] Klimczak, M. *Poincaré duality for spaces with isolated singularities*, Manuscripta Mathematica, **153** (2017), pp. 231–262 DOI: [10.1007/s00229-016-0884-5](https://doi.org/10.1007/s00229-016-0884-5)
- [27] McConnell, M. *The rational homology of toric varieties is not a combinatorial invariant*, Proceedings of the American Mathematical Society, **105** (1989), pp. 986–991 DOI: [10.2307/2047063](https://doi.org/10.2307/2047063)
- [28] Rourke, C., Sanderson, B., *Homology stratifications and intersection homology*, Geometry & Topology Monographs, **2** (1999), pp. 455–472 DOI: [10.2140/gtm.1999.2.455](https://doi.org/10.2140/gtm.1999.2.455)
- [29] Schlöder, F. & Essig, J. T. *Multiplicative de Rham Theorems for Relative and Intersection Space Cohomology*, J. of Singularities, **19** (2019), pp. 97–130 DOI: [10.5427/jsing.2019.19g](https://doi.org/10.5427/jsing.2019.19g)
- [30] Sella, Y. *Comparison of sheaf cohomology and singular cohomology*, (2016) arXiv: [1602.06674](https://arxiv.org/abs/1602.06674)
- [31] Spiegel, M. *K-theory of intersection spaces*, PhD thesis (Universität Heidelberg 2013)
- [32] Stanley, R. *Generalized H-vectors, intersection cohomology of toric varieties, and related results*, Commutative Algebra And Combinatorics, **11** (1987), pp. 187–214
- [33] Wrazidlo, D. *A fundamental class for intersection spaces of depth one Witt spaces*, Manuscripta Mathematica, **166** (2021), pp. 199–236 DOI: [10.1007/s00229-020-01238-7](https://doi.org/10.1007/s00229-020-01238-7)
- [34] Wrazidlo, D. *Induced maps between intersection spaces*, Master thesis, (Universität Heidelberg, 2013)
- [35] Yavin, D. *The intersection homology with twisted coefficients of toric varieties*, PhD thesis, (Massachusetts Institute of Technology, 1990)

MARKUS BANAGL, INSTITUT FÜR MATHEMATIK, UNIVERSITÄT HEIDELBERG, IM NEUENHEIMER FELD 205, 69120 HEIDELBERG, GERMANY

Email address: banagl@mathi.uni-heidelberg.de

SHAHRYAR GHAED SHARAF, FAKULTÄT FÜR MATHEMATIK UND INFORMATIK, FRIEDRICH-SCHILLER-UNIVERSITÄT JENA, ERNST-ABBE-PLATZ 2 , 07743 JENA, GERMANY

Email address: shahryar.ghaed.sharaf@uni-jena.de